

PATENT COOPERATION TREATY

20 SEP 2004

PCT

From the INTERNATIONAL BUREAU

NOTIFICATION OF RECEIPT OF
RECORD COPY

(PCT Rule 24.2(a))

To:

MEYERS, Hans-Wilhelm
Postfach 10 22 41
50462 Köln
Germany

AvX Sg W Da Hi HPJ ME TW JH KB

19. MAI 2003

CS K 7 21-0003

Date of mailing (day/month/year) 28 April 2003 (28.04.03)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference 030640woMebS	International application No. PCT/EP03/02857

The applicant is hereby notified that the International Bureau has received the record copy of the international application as detailed below.

Name(s) of the applicant(s) and State(s) for which they are applicants:

EVOTEC NEUROSCIENCES GMBH (for all designated States except US)

VON DER KAMMER, Heinz et al (for US)

International filing date : 19 March 2003 (19.03.03)

Priority date(s) claimed : 21 March 2002 (21.03.02)

21 March 2002 (21.03.02)

Date of receipt of the record copy
by the International Bureau : 15 April 2003 (15.04.03)

List of designated Offices :

AP : GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW

EA : AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

EP : AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR

OA : BF, BJ, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG

National : AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ,
EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Authorized officer:

R. Grenier (Fax 338 8970)

Facsimile No. (41-22) 338.89.70

Telephone No. (41-22) 338 8277

PRIORITY COOPERATION TREATY

PCT

NOTIFICATION CONCERNING
SUBMISSION OR TRANSMITTAL
OF PRIORITY DOCUMENT

(PCT Administrative Instructions, Section 411)

From the INTERNATIONAL BUREAU

To:

MEYERS, Hans-Wilhelm
Postfach 10 22 41
50462 Köln
Germany

AV	K	G	W	Da	Hi	HP	ME	TW	JH	KE
06.0KT.2003										
CS K F 21.10.03										

Date of mailing (day/month/year) 30 September 2003 (30.09.03)	
Applicant's or agent's file reference 030640woMebs	IMPORTANT NOTIFICATION
International application No. PCT/EP03/02857	International filing date (day/month/year) 19 March 2003 (19.03.03)
International publication date (day/month/year) Not yet published	Priority date (day/month/year) 21 March 2002 (21.03.02)
Applicant EVOTEC NEUROSCIENCES GMBH et al	

- The applicant is hereby notified of the date of receipt (except where the letters "NR" appear in the right-hand column) by the International Bureau of the priority document(s) relating to the earlier application(s) indicated below. Unless otherwise indicated by an asterisk appearing next to a date of receipt, or by the letters "NR", in the right-hand column, the priority document concerned was submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b).
- This updates and replaces any previously issued notification concerning submission or transmittal of priority documents.
- An asterisk(*) appearing next to a date of receipt, in the right-hand column, denotes a priority document submitted or transmitted to the International Bureau but not in compliance with Rule 17.1(a) or (b). In such a case, **the attention of the applicant is directed** to Rule 17.1(c) which provides that no designated Office may disregard the priority claim concerned before giving the applicant an opportunity, upon entry into the national phase, to furnish the priority document within a time limit which is reasonable under the circumstances.
- The letters "NR" appearing in the right-hand column denote a priority document which was not received by the International Bureau or which the applicant did not request the receiving Office to prepare and transmit to the International Bureau, as provided by Rule 17.1(a) or (b), respectively. In such a case, **the attention of the applicant is directed** to Rule 17.1(c) which provides that no designated Office may disregard the priority claim concerned before giving the applicant an opportunity, upon entry into the national phase, to furnish the priority document within a time limit which is reasonable under the circumstances.

<u>Priority date</u>	<u>Priority application No.</u>	<u>Country or regional Office or PCT receiving Office</u>	<u>Date of receipt of priority document</u>
21 Marc 2002 (21.03.02)	02006353.3	EP	16 Sept 2003 (16.09.03)
21 Marc 2002 (21.03.02)	60/365,815	US	12 June 2003 (12.06.03)

OK
09.10.03

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No. (41-22) 338.89.70	Authorized officer Elisabeth KÖNIG (Fax 338 8970) Telephone No. (41-22) 338 8748
--	--

PATENT COOPERATION TREATY

From the INTERNATIONAL BUREAU

PCT

NOTICE INFORMING THE APPLICANT OF THE
COMMUNICATION OF THE INTERNATIONAL
APPLICATION TO THE DESIGNATED OFFICES

(PCT Rule 47.1(c), first sentence)

To:

20 SEP 2004

MEYERS, Hans-Wilhelm
Postfach 10 22 41
50462 Köln
ALLEMAGNE

K Sg W Da Hi HP ME TW JH KB

13. OKT. 2003

CS K 7 21. 10. 03

Date of mailing(day/month/year)
02 October 2003 (02.10.03)Applicant's or agent's file reference
030640woMebs

IMPORTANT NOTICE

International application No.
PCT/EP03/02857International filing date(day/month/year)
19 March 2003 (19.03.03)Priority date(day/month/year)
21 March 2002 (21.03.02)

Applicant

EVOTEC NEUROSCIENCES GMBH

1. Notice is hereby given that the International Bureau has **communicated**, as provided in Article 20, the international application to the following designated Offices on the date indicated above as the date of mailing of this notice:

AU, AZ, BY, CH, CN, CO, DE, DZ, HU, JP, KG, KP, KR, MD, MK, MZ, RU, TM, US

In accordance with Rule 47.1(c), third sentence, those Offices will accept the present notice as conclusive evidence that the communication of the international application has duly taken place on the date of mailing indicated above and no copy of the international application is required to be furnished by the applicant to the designated Office(s).

2. The following designated Offices have waived the requirement for such a communication at this time:

AE, AG, AL, AM, AP, AT, BA, BB, BG, BR, BZ, CA, CR, CU, CZ, DK, DM, EA, EC, EE, EP, ES, FI, GB, GD, GE, GH, GM, HR, ID, IL, IN, IS, KE, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MG, MN, MW, MX, NI, NO, NZ, OA, OM, PH, PL, PT, RO, SC, SD, SE, SG, SK, SL, TJ, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW

The communication will be made to those Offices only upon their request. Furthermore, those Offices do not require the applicant to furnish a copy of the international application (Rule 49.1(a-bis)).

3. Enclosed with this notice is a copy of the international application as published by the International Bureau on 02 October 2003 (02.10.03) under No. 03/080661

4. **TIME LIMITS for filing a demand for international preliminary examination and for entry into the national phase**

The applicable time limit for entering the national phase will, **subject to what is said in the following paragraph**, be **30 MONTHS** from the priority date, not only in respect of any elected Office if a demand for international preliminary examination is filed before the expiration of **19 months** from the priority date, but also in respect of any designated Office, in the absence of filing of such demand, where Article 22(1) as modified with effect from 1 April 2002 applies in respect of that designated Office. For further details, see *PCT Gazette* No. 44/2001 of 1 November 2001, pages 19926, 19932 and 19934, as well as the *PCT Newsletter*, October and November 2001 and February 2002 issues.

In practice, **time limits other than the 30-month time limit** will continue to apply, for various periods of time, in respect of certain designated or elected Offices. For **regular updates on the applicable time limits** (20, 21, 30 or 31 months, or other time limit), Office by Office, refer to the *PCT Gazette*, the *PCT Newsletter* and the *PCT Applicant's Guide*, Volume II, National Chapters, all available from WIPO's Internet site, at <http://www.wipo.int/pc/en/index.html>.

For filing a **demand for international preliminary examination**, see the *PCT Applicant's Guide*, Volume I/A, Chapter IX. Only an applicant who is a national or resident of a PCT Contracting State which is bound by Chapter II has the right to file a demand for international preliminary examination (at present, all PCT Contracting States are bound by Chapter II).

It is the applicant's **sole responsibility** to monitor all these time limits.

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Authorized officer

Judith Zahra

Facsimile No.(41-22) 740.14.35

Telephone No.(41-22) 338.91.11

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

20 SEP 2004

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference 030640woMebS	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/EP 03/ 02857	International filing date (day/month/year) 19/03/2003	(Earliest) Priority Date (day/month/year) 21/03/2002
Applicant EVOTEC NEUROSCIENCES GMBH		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :

☒ contained in the international application in written form.

☒ filed together with the international application in computer readable form.

☐ furnished subsequently to this Authority in written form.

☐ furnished subsequently to this Authority in computer readable form.

☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☒ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the title,

☐ the text is approved as submitted by the applicant.

☒ the text has been established by this Authority to read as follows:

DIAGNOSTIC AND THERAPEUTIC USE OF HUMAN MAGUIN PROTEINS AND NUCLEIC ACIDS FOR NEURODEGENERATIVE DISEASE

5. With regard to the abstract,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the drawings to be published with the abstract is Figure No.

☒ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

1
☐ None of the figures.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP/02857

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/47 C12N15/12 C12N5/10 C07K16/18 G01N33/53
C12Q1/68 A61K38/16 A61K48/00 A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K G01N C12Q A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

GENSEQ, EMBL, EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL 'Online! 16 January 2002 (2002-01-16) LANIGAN, T.M. AND GUAN, K.L.: "Homo sapiens connector enhancer of KSR2A mRNA, complete cds." Database accession no. AF418269 XP002210679 --- -/--	11, 13, 14, 16, 20, 26, 28, 29



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

18 June 2003

Date of mailing of the international search report

30/06/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

ALCONADA RODRIG..., A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP/02857

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE GENSEQ 'Online! 8 November 2000 (2000-11-08) TAKAI, Y. ET AL.: "Rat MAGUIN 1 protein" Database accession no. AAY92942 XP002210736 -& DATABASE WPI Section Ch, Week 200033 Derwent Publications Ltd., London, GB; Class B04, AN 2000-387785 XP002210680 & WO 00 29572 A (KAGAKU GIJUTSU SHINKO JIGYODAN), 25 May 2000 (2000-05-25) abstract</p> <p>---</p>	11,13, 14,16, 20,26, 28,29
X	<p>DATABASE GENSEQ 'Online! 8 November 2000 (2000-11-08) TAKAI, Y. ET AL.: "Rat MAGUIN 2 protein" Database accession no. AAY92943 XP002210737 -& DATABASE WPI Section Ch, Week 200033 Derwent Publications Ltd., London, GB; Class B04, AN 2000-387785 XP002210680 & WO 00 29572 A (KAGAKU GIJUTSU SHINKO JIGYODAN), 25 May 2000 (2000-05-25) abstract</p> <p>-----</p>	11,13, 14,16, 20,26, 28,29

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 12-15 (in part) and 19, 21-25 (complete)

Present claim 12 relates to a method of treating or preventing a neurodegenerative disease, said method comprising administering to said subject an amount of an agent, wherein said agent is defined by reference to a desirable characteristic or property, namely, that said agent directly or indirectly modulates the activity or level of the (i) gene coding for human MAGUIN-1 and/or human MAGUIN-2; and/or (ii) a transcription product of a gene coding for human MAGUIN-1 or human MAGUIN-2 and/or (iii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2 and/or a fragment, or derivative, or variant of (i) to (iii). The claims cover the methods that involve the administration of an agent having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT for only a very limited number of such agents. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the agent by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the methods which make use of the agent, wherein the agent comprises the human MAGUIN-1 or MAGUIN-2 polypeptide sequences or the corresponding polynucleotide sequence in sense or antisense orientation (see page 14 from the description).

The same objection applies to claim 13, which refers to the above mentioned agent or modulator as such, to claim 14 which refers to pharmaceutical compositions comprising said modulator and to claim 15, which refers to the second medical use of said modulator.

Present claim 19 relates to a method for the identification of a compound for inhibition of binding between a ligand and human MAGUIN-1 or MAGUIN-2; wherein said method implies contacting MAGUIN-1 or MAGUIN-2 with a detectable ligand in the presence of the compound to be tested. The claim covers all possible MAGUIN ligands, whereas the application does not provide support within the meaning of Article 6 PCT or disclosure within the meaning of Article 5 PCT for any of such MAGUIN ligands. In the present case, the claim so lacks support, and the application so lacks disclosure, that a meaningful search of the claim is impossible. Independent of the above reasoning, the claim also lacks clarity (Article 6 PCT). An attempt is made to define the product by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been not carried out for those claims.

Claims 21-25 relate to methods of producing medicaments and medicaments

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

obtained by said method, wherein said medicaments contain a compound which is defined by reference to a desirable characteristic or property, namely, that the compound can be identified by the methods of claim 18-20. The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT for none of such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search of the claim impossible. Consequently, the search has not been carried out for those claims.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International application No.
EP 03/02857

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 12 and 17 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 12-15 (in part) and 19, 21-25 (complete)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

PATENT COOPERATION TREATY

PCT

20 SEP 2004

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

(Rationalised Report according to the Notice of the President of the EPO published in the OJ11/2001)

Applicant's or agent's file reference 030640wo/Me/sto	<div style="display: flex; justify-content: space-between;"> FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416) </div>	
International application No. PCT/EP03/02857	International filing date (day/month/year) 19/03/2003	Priority date (day/month/year) 21/03/2002
International Patent Classification (IPC) or national classification and IPC C07K14/47		
Applicant EVOTEC NEUROSCIENCES GMBH ET AL.		


1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2. This **REPORT** consists of a total of 2 sheets, including this cover sheet.

☐ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consists of a total of _____ sheets.

3. This report contains indications relating to the following items:
 - I ☒ Basis of the report
 - II ☐ Priority
 - III ☒ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
 - IV ☐ Lack of unity of invention
 - V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
 - VI ☐ Certain documents cited
 - VII ☐ Certain defects in the international application
 - VIII ☐ Certain observations on the international application

Date of submission of the demand 20/10/2003	Date of completion of this report 16/02/2004
Name and mailing address of the IPEA/ European Patent Office, Gitschiner Str. 103 D-10969 Berlin - Germany Tel.: (+49-30) 25901-0 Fax: (+49-30) 25901-840	Authorized officer DE BUNDEL E R J Tel. (+49-89) 2399 2828



I. Basis of the report

The basis of this international preliminary examination is the application as originally filed.

III. Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

The question of whether the claimed invention appears to be novel, to involve an inventive step, or to be industrially applicable has not been the subject of the international preliminary examination in respect of the claims which have not been searched (Article 17(2)(a) or (3) and Rule 66.1(e) PCT); see also international search report).

V. Reasoned statement under Rule 66.2(a)(ii) with regard to novelty, inventive step or industrial applicability

To the extent that the international preliminary examination has been carried out (see item III above), the following is pointed out:

In light of the documents cited in the international search report, it is considered that the invention as defined in at least some of the claims, which have been the subject of an international search report, does not appear to meet the criteria mentioned in Article 33(1) PCT, i.e. does not appear to be novel and/or to involve an inventive step (see international search report, in particular the documents cited X and/or Y and corresponding claim references).

(19) World Intellectual Property Organization
International Bureau



20 SEP 2004



(43) International Publication Date
2 October 2003 (02.10.2003)

PCT

(10) International Publication Number
WO 03/080661 A1

(51) International Patent Classification⁷: **C07K 14/47**,
C12N 15/12, 5/10, C07K 16/18, G01N 33/53, C12Q 1/68,
A61K 38/16, 48/00, A01K 67/027

(21) International Application Number: PCT/EP03/02857

(22) International Filing Date: 19 March 2003 (19.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
02006353.3 21 March 2002 (21.03.2002) EP
60/365,815 21 March 2002 (21.03.2002) US

(71) Applicant (for all designated States except US): **EVOTEC
NEUROSCIENCES GMBH** [DE/DE]; Schnackenburg-
allee 114, 22525 Hamburg (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **VON DER KAM-
MER, Heinz** [DE/DE]; Verbindungsstr. 6d, 22607 Ham-
burg (DE). **HANES, Jozef** [SK/DE]; Dornkamp 15, 22869

Schenefeld (DE). **POHLNER, Johannes** [DE/DE]; Quit-
tenweg 11, 22175 Hamburg (DE).

(74) Agents: **MEYERS, Hans-Wilhelm** et al.; Postfach 10 22
41, 50462 Köln (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD,
SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report

[Continued on next page]

(54) Title: **DIAGNOSTIC AND THERAPEUTIC USE OF HUMAN MAGUIN PROTEINS AND NUCLEIC ACIDS FOR NEU-
RODEGENERATIVE DISEASES**

Identification of Genes Involved in Alzheimer's Disease Pathology



(57) Abstract: The present invention discloses the differential expression of the human MAGUIN-1 caveolae-associated integral membrane protein flotillin-1 and/or human MAGUIN-2 gene in specific brain regions of Alzheimer's disease patients. Based on this finding, this invention provides a method for diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, or other neurodegenerative diseases in a subject patient, or for determining whether a subject is at increased risk of developing such a disease or other related neurodegenerative diseases. Furthermore, this invention provides therapeutic and prophylactic methods for treating or preventing Alzheimer's disease and related neurodegenerative disorders using a human MAGUIN caveolae-associated integral membrane protein gene. A method of screening for modulating agents of neurodegenerative diseases is also disclosed.

Best Available Copy



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

DIAGNOSTIC AND THERAPEUTIC USE OF HUMAN MAGUIN PROTEINS AND NUCLEIC ACIDS FOR NEURODEGENERATIVE DISEASES

The present invention relates to methods of diagnosing, prognosticating and monitoring the progression of neurodegenerative diseases in a subject. Furthermore, methods of therapy control and screening for modulating agents of neurodegenerative diseases are provided. The invention also discloses pharmaceutical compositions, kits, and recombinant animal models.

Neurodegenerative diseases, in particular Alzheimer's disease (AD), have a strongly debilitating impact on a patient's life. Furthermore, these diseases constitute an enormous health, social, and economic burden. AD is the most common neurodegenerative disease, accounting for about 70% of all dementia cases, and it is probably the most devastating age-related neurodegenerative condition affecting about 10% of the population over 65 years of age and up to 45% over age 85 (for a recent review see Vickers et al., *Progress in Neurobiology* 2000, 60: 139-165). Presently, this amounts to an estimated 12 million cases in the US, Europe, and Japan. This situation will inevitably worsen with the demographic increase in the number of old people ("aging of the baby boomers") in developed countries. The neuropathological hallmarks that occur in the brains of individuals with AD are senile plaques, composed of amyloid- β protein, and profound cytoskeletal changes coinciding with the appearance of abnormal filamentous structures and the formation of neurofibrillary tangles.

The amyloid- β (A β) protein evolves from the cleavage of the amyloid precursor protein (APP) by different kinds of proteases. The cleavage by the β/γ -secretase leads to the formation of A β peptides of different lengths, typically a short more soluble and slow aggregating peptide consisting of 40 amino acids and a longer 42 amino acid peptide, which rapidly aggregates outside the cells, forming the characteristic amyloid plaques (Selkoe, *Physiological Rev* 2001, 81: 741-66; Greenfield et al., *Frontiers Bioscience* 2000, 5: D72-83). Two types of plaques, diffuse plaques and neuritic plaques, can be detected in the brain of AD patients, the latter ones being the classical, most prevalent type. They are primarily found in the cerebral cortex and hippocampus. The neuritic plaques have a diameter of 50 μ m to 200 μ m and are composed of insoluble fibrillar amyloids, fragments of dead neurons, of microglia and astrocytes, and other components such as neurotransmitters, apolipoprotein E, glycosaminoglycans, α 1-antichymotrypsin and others. The generation of toxic A β deposits in the brain starts very early in the course of AD, and it is discussed to be a key player for the subsequent

destructive processes leading to AD pathology. The other pathological hallmarks of AD are neurofibrillary tangles (NFTs) and abnormal neurites, described as neuropil threads (Braak and Braak, *Acta Neuropathol* 1991, 82: 239-259). NFTs emerge inside neurons and consist of chemically altered tau, which forms paired helical filaments twisted around each other. Along the formation of NFTs, a loss of neurons can be observed. It is discussed that said neuron loss may be due to a damaged microtubule-associated transport system (Johnson and Jenkins, *J Alzheimers Dis* 1996, 1: 38-58; Johnson and Hartigan, *J Alzheimers Dis* 1999, 1: 329-351). The appearance of neurofibrillary tangles and their increasing number correlates well with the clinical severity of AD (Schmitt et al., *Neurology* 2000, 55: 370-376).

AD is a progressive disease that is associated with early deficits in memory formation and ultimately leads to the complete erosion of higher cognitive function. The cognitive disturbances include among other things memory impairment, aphasia, agnosia and the loss of executive functioning. A characteristic feature of the pathogenesis of AD is the selective vulnerability of particular brain regions and subpopulations of nerve cells to the degenerative process. Specifically, the temporal lobe region and the hippocampus are affected early and more severely during the progression of the disease. On the other hand, neurons within the frontal cortex, occipital cortex, and the cerebellum remain largely intact and are protected from neurodegeneration (Terry et al., *Annals of Neurology* 1981, 10: 184-92).

The age of onset of AD may vary within a range of 50 years, with early-onset AD occurring in people younger than 65 years of age, and late-onset of AD occurring in those older than 65 years. About 10% of all AD cases suffer from early-onset AD, with only 1-2% being familial, inherited cases.

Currently, there is no cure for AD, nor is there an effective treatment to halt the progression of AD or even to diagnose AD ante-mortem with high probability. Several risk factors have been identified that predispose an individual to develop AD, among them most prominently the epsilon 4 allele of the three different existing alleles (epsilon 2, 3, and 4) of the apolipoprotein E gene (ApoE) (Strittmatter et al., *Proc Natl Acad Sci USA* 1993, 90: 1977-81; Roses, *Ann NY Acad Sci* 1998, 855: 738-43). The polymorphic plasmaprotein ApoE plays a role in the intercellular cholesterol and phospholipid transport by binding low-density lipoprotein receptors, and it seems to play a role in neurite growth and regeneration. Efforts to detect further susceptibility genes and disease-linked polymorphisms, lead to the assumption that specific regions and genes on human chromosomes 10 and 12 may be associated with late-onset AD (Myers et al., *Science* 2000, 290: 2304-5; Bertram et al., *Science* 2000, 290: 2303; Scott et al., *Am J Hum Genet* 2000, 66: 922-32).

Although there are rare examples of early-onset AD which have been attributed to genetic defects in the genes for amyloid precursor protein (APP) on chromosome 21, presenilin-1 on chromosome 14, and presenilin-2 on chromosome 1, the prevalent form of late-onset sporadic AD is of hitherto unknown etiologic origin. The mutations found to date account for only half of the familial AD cases, which is less than 2% of all AD patients. The late onset and complex pathogenesis of neurodegenerative disorders pose a formidable challenge to the development of therapeutic and diagnostic agents. It is pivotal to expand the pool of potential drug targets and diagnostic markers. It is therefore an object of the present invention to provide insight into the pathogenesis of neurological diseases and to provide methods, materials, agents, compositions, and animal models which are suited inter alia for the diagnosis and development of a treatment of these diseases. This object has been solved by the features of the independent claims. The subclaims define preferred embodiments of the present invention.

Neurons conduct signals in the form of electrical impulses. The communication between the cells of a neuronal network involves several chemical steps, including the production, release, and transport of signal molecules and the recognition of such messengers by a receptor. These steps take place at the anatomical contact of an axon with the dendrites (axonodendritic), the cell body (axonosomatic), or rarely with the axon of another neuron, and also between neurons and cells of muscle- and gland tissue. Trillions of such specialized cell junctions (synapses) in the human brain are crucial for controlling mental activity and learning processes (Tessier-Lavigne and Goodman, *Science* 1996, 274: 1123-1133). A synapse is composed of the presynaptic element, the synaptic cleft with a spacing distance of about 20-30 nm and the postsynaptic component. They are responsible for altering the membrane potential of the postsynaptic neuron or other effector cells. The quality and intensity of information transferred relies on the number, location, and distribution of synapses. The basis of learning and memory is believed to be due to brain plasticity, i.e. to the plasticity of synapses. The storage and processing of information cause alterations in the structure, chemistry, and strength of synapses and the formation of new synapses (Poirazi and Mel, *Neuron* 2001, 29: 779-796). The postsynaptic density (PSD) is a specialized synaptic signaling assemblage, composed of a specific set of proteins which are assembled together and linked to the cytoplasmic face of the postsynaptic membrane. An important function of the postsynaptic density is the provision of a structural matrix consisting of cytoskeletal and regulatory proteins which localize and accumulate neurotransmitter receptors (e.g. glutamate receptors) and anchor signaling molecules at the postsynaptic membrane (Sheng and Kim, *Current Opinion Neurobiology* 1996, 6:

602-608). Neurotransmitter receptors convert the extracellular chemical signals into intracellular signals. Neurotransmitters are released from the presynaptic membrane into the synaptic cleft via exocytotic processes (vesicle formation). At the postsynaptic membrane they are bound by their specific receptors, resulting in the generation of second messengers. Thus, neurotransmitters function as mediators of nerve impulse transmission across the synapse. The PSD organizes and regulates postsynaptic signal transduction (Kim and Huganir, *Current Opinion Cell Biology* 1999, 11:248-254). To date, several components of the PSD have been identified, among them the prototypic synaptic scaffolding protein postsynaptic density (PSD)-95/synapse-associated protein (SAP) 90. PSD-95/SAP90 and its isoforms belong to a family of membrane-associated guanylate kinases (MAGUK) (Hirao et al., *Journal of Biological Chemistry* 1998, 273: 21105-21110; Hirao et al., *Journal of Biological Chemistry* 2000, 275: 2966-2972). The members of the MAGUK protein family are multiple PDZ-domain containing proteins which interact with many neuronal adhesion proteins, receptors (e.g. N-methyl-D-aspartate (NMDA)) and ion-channels (e.g. potassium channels) through these domains and mediate their assembling at the PSD (Kim et al., *Neuron* 1996, 17: 103-113).

In this context, a novel neuronal membrane-associated guanylate kinase-interacting protein, denoted MAGUIN, was found by Yao et al. (*Journal of Biological Chemistry* 1999, 274: 11889-11896). Using the PDZ-domain sequence of the neurospecific synaptic scaffolding molecule (S-SCAM) as bait, the authors screened a rat brain yeast two-hybrid library and obtained several positive clones, among them two clones subsequently named rat MAGUIN-1 and rat MAGUIN-2 (GenBank accession numbers AF102853 and AF102854, respectively). A Northern blot analysis of different rat tissues with a rat MAGUIN-1 probe revealed brain specific hybridization signals at 4.4 kDa and 5.4 kDa. MAGUIN proteins have a chimerical molecular structure consisting of several protein modules, an N-terminally located sterile alpha-motif (SAM) (aa 8-75), a PDZ-domain (aa 156-296) and a C-terminally located Pleckstrin-homology (PH)-domain (aa 571-667). The Sterile Alpha Motif (SAM) contains four different domain structures, spanning approximately 70 amino acids generating a compact five-helix bundle with a highly conserved hydrophobic core. This motif was found to be part of a number of types of proteins, including signal transduction proteins, playing a role in mediating protein-protein interactions or DNA binding (Schultz et al., *Protein Science* 1997, 6: 249-253). The term PDZ is named after three proteins (i.e. PSD-95/SAP90, *Drosophila* discs-large tumor suppressor protein (Dlg-A) and the tight junction protein ZO-1), originally identified as proteins sharing the same repeats of about 90 amino acids. These amino acids form a distinctive structure of two α helices and six β sheets which mediate the interaction with the carboxyl termini of various proteins. Often, PDZ-

domains of different proteins can heterodimerize with each other. The target proteins are transmembrane receptors, ion channels, or signaling proteins. PDZ-domain binding seems to be important in receptor clustering and in recruiting signal transduction molecules to the plasma membrane. PDZ-domain harboring proteins are for example tyrosine phosphatases and the previously described membrane-associated guanylate kinase-like proteins (Doyle et al., *Cell* 1996, 85: 1067-1076). The Pleckstrin-homology (PH)-domain forms an anti-parallel perpendicular β -sheet sandwich with a succeeding α -helical structure. The β -sheet-loops are important for high affinity binding to specific phosphatidylinositol phosphates, allowing signaling proteins containing the PH-domain to anchor to membranes, for example as studied for GTP binding proteins, protein kinases and phospholipase C isoforms (Lemmon et al., *Trends Cell Biology* 1997, 7: 237-242).

Rat MAGUIN-1 and rat MAGUIN-2 cover 3099 and 2691 nucleotides of coding sequence, respectively, (GenBank accession numbers AF102853, AF102854) which encode for proteins of 1032 aa amino acids (rat MAGUIN-1) (GenBank accession number aad04568) and 896 amino acids (rat MAGUIN-2) (GenBank accession number aad04567), respectively. Rat MAGUIN-2 lacks the 3' terminal PDZ binding motif of rat MAGUIN-1. In neurons, the rat MAGUIN proteins are localized in the cell body and in neurites where they are associated with the plasma-membrane via their PH-domains. Full-length rat MAGUIN-1 could be recovered from neuronal membrane fractions. Rat MAGUIN-1, but not rat MAGUIN-2, interacts via the PDZ-binding motif with the PDZ-domain of PSD-95/SAP90 and the synaptic scaffolding molecule (S-SCAM) (Hirao et al., *Journal of Biological Chemistry* 1998, 273: 21105-21110). S-SCAM has recently been identified as a multiple PDZ-domain containing protein, interacting with SAP90/PSD-95-associated protein (SAPAP), the NMDA receptor, and neuronal adhesion molecules. Additionally, interaction of the PH-domain of rat MAGUIN-1 with the kinase domain of Raf could be confirmed *in vitro* and *in vivo*, but there is no evidence for the activation of Raf or its recruitment to the plasma membrane by rat MAGUIN-1.

To date, the precise function of rat MAGUIN-1 and rat MAGUIN-2 is still not clear. Some clues for the role of rat MAGUIN-1 in the cellular context come from particular protein domains and motifs, specific brain expression patterns, and structural homologies to other proteins like CNK (connector enhancer of kinase suppressor of ras (ksr)), as described for the fruit fly. CNK binds kinase suppressor for Ras and Raf kinase, functions in the Ras/MAP kinase pathway, and has been found to play a profound role in the regulation of eye development (Therrien et al., *Cell* 1998, 95: 343-353). On the basis of the similarities between *Drosophila* CNK (contains SAM, PDZ and PH-domains) and rat MAGUIN-1, rat MAGUIN-1 is discussed as a rodent homolog to CNK. Thus, it is

likely that rat MAGUIN-1 assembles components of synaptic junctions and regulators of the MAP kinase pathway and links them to NMDA receptors and neuronal adhesion molecules. To date, no experiments have been described that show a relationship between a differential expression of human MAGUIN genes, neither on a transcriptional nor on a translational level, and the pathology of neurodegenerative disorders.

The disclosure in the present invention of the human MAGUIN-1 gene and the identification of a link of both human MAGUIN-1 and/or human MAGUIN-2 to neurodegenerative diseases, particularly Alzheimer's disease, offers new ways, inter alia, for the diagnosis and treatment of such diseases.

The singular forms "a", "an", and "the" as used herein and in the claims include plural reference unless the context dictates otherwise. For example "a cell" means as well a plurality of cells, and so forth. The term "and/or" as used in the present specification and in the claims implies that the phrases before and after this term are to be considered either as alternatives or in combination. For instance, the wording "determination of a level and/or an activity" means that either only a level, or only an activity, or both a level and an activity are determined. The term "level" as used herein is meant to comprise a gage of, or a measure of the amount of, or a concentration of a transcription product, for instance an mRNA, or a translation product, for instance a protein or polypeptide. The term "activity" as used herein shall be understood as a measure for the ability of a transcription product or a translation product to produce a biological effect or a measure for a level of biologically active molecules. The term "activity" also refers to enzymatic activity. The terms "level" and/or "activity" further refer to gene expression levels or gene activity. Gene expression can be defined as the utilization of the information contained in a gene by transcription and translation leading to the production of a gene product. "Dysregulation" shall mean an upregulation or downregulation of gene expression. A gene product comprises either RNA or protein and is the result of expression of a gene. The amount of a gene product can be used to measure how active a gene is. The term "gene" as used in the present specification and in the claims comprises both coding regions (exons) as well as non-coding regions (e.g. non-coding regulatory elements such as promoters or enhancers, introns, leader and trailer sequences). The term "ORF" is an acronym for "open reading frame" and refers to a nucleic acid sequence that does not possess a stop codon in at least one reading frame and therefore can potentially be translated into a sequence of amino acids. "Regulatory elements" as used in the present disclosure shall comprise inducible and non-inducible promoters, enhancers, operators and other elements that drive and regulate gene expression. The term "fragment" as used herein is meant to comprise e.g.

an alternatively spliced, or truncated, or otherwise cleaved transcription product or translation product. The term "derivative" as used herein refers to a mutant, or an RNA-edited, or a chemically modified, or otherwise altered transcription product, or to a mutant, or chemically modified, or otherwise altered translation product. For instance, a "derivative" may be generated by processes such as altered phosphorylation, or glycosylation, or acetylation, or lipidation, or by altered signal peptide cleavage or other types of maturation cleavage. These processes may occur post-translationally. The term "modulator" as used in the present invention and in the claims refers to a molecule capable of changing or altering the level and/or the activity of a gene, or a transcription product of a gene, or a translation product of a gene. Preferably, a "modulator" is capable of changing or altering the biological activity of a transcription product or a translation product of a gene. Said modulation, for instance, may be an increase or a decrease in enzyme activity, a change in binding characteristics, or any other change or alteration in the biological, functional, or immunological properties of said translation product of a gene.

The terms "agent", "reagent", or "compound" refer to any substance, chemical, composition or extract that have a positive or negative biological effect on a cell, tissue, body fluid, or within the context of any biological system, or any assay system examined. They can be agonists, antagonists, partial agonists or inverse agonists of a target. Such agents, reagents, or compounds may be nucleic acids, natural or synthetic peptides or protein complexes, or fusion proteins. They may also be antibodies, organic or anorganic molecules or compositions, small molecules, drugs and any combinations of any of said agents above. They may be used for testing, for diagnostic or for therapeutic purposes. The terms "oligonucleotide primer" or "primer" refer to short nucleic acid sequences which can anneal to a given target polynucleotide by hybridization of the complementary base pairs and can be extended by a polymerase. They may be chosen to be specific to a particular sequence or they may be randomly selected, e.g. they will prime all possible sequences in a mix. The length of primers used herein may vary from 10 nucleotides to 80 nucleotides. "Probes" are short nucleic acid sequences of the nucleic acid sequences described and disclosed herein or sequences complementary therewith. They may comprise full length sequences, or fragments, derivatives, isoforms, or variants of a given sequence. The identification of hybridization complexes between a "probe" and an assayed sample allows the detection of the presence of other similar sequences within that sample. As used herein, "homolog or homology" is a term used in the art to describe the relatedness of a nucleotide or peptide sequence to another nucleotide or peptide sequence, which is determined by the degree of identity and/or similarity between said sequences compared.

The term "variant" as used herein refers to any polypeptide and protein, in reference to polypeptides and proteins disclosed in the present invention, in which one or more amino acids are added and/or substituted and/or deleted and/or inserted at the N-terminus, and/or the C-terminus, and/or within the native amino acid sequences of the native polypeptides or proteins of the present invention. Furthermore, the term "variant" shall include any shorter or longer version of a polypeptide or protein. "Variants" shall also comprise a sequence that has at least about 80% sequence identity, more preferably at least about 90% sequence identity, and most preferably at least about 95% sequence identity with the amino acid sequences of SEQ ID NO. 1 and/or SEQ ID NO. 2, respectively. Derivatives, variants, and fragments may include, but are not limited to, a functional SAM, a functional PDZ and a functional PH domain or other functional modules within the polypeptide sequence of human MAGUIN-1 and/or human Maguin-2 proteins. "Variants" of a protein molecule shown in SEQ ID NO. 1 and/or SEQ ID NO. 2 may include, for example, proteins with conservative amino acid substitutions in highly conservative regions. For example, isoleucine, valine and leucine can each be substituted for one another. Aspartate and glutamate can be substituted for each other. Glutamine and asparagine can be substituted for each other. Serine and threonine can be substituted for each other. Amino acid substitutions in less conservative regions include, for example, isoleucine, valine and leucine, which can each be substituted for one another. Aspartate and glutamate can be substituted for each other. Glutamine and asparagine can be substituted for each other. Serine and threonine can be substituted for each other. Glycine and alanine can be substituted for each other. Alanine and valine can be substituted for each other. Methionine can be substituted for each of leucine, isoleucine or valine, and vice versa. Lysine and arginine can be substituted for each other. One of aspartate and glutamate can be substituted for one of arginine or lysine, and vice versa. Histidine can be substituted for arginine or lysine, and vice versa. Glutamine and glutamate can be substituted for each other. Asparagine and aspartate can be substituted for each other. Other examples of protein modifications include glycosylation and further post-translational modifications. "Proteins and polypeptides" of the present invention include variants, fragments and chemical derivatives of the protein comprising SEQ ID NO. 1 and/or SEQ ID NO. 2. They can include proteins and polypeptides which can be isolated from nature or which can be produced by recombinant and/or synthetic means. Native proteins or polypeptides refer to naturally-occurring truncated or secreted forms, naturally occurring variant forms (e.g. splice variants) and naturally occurring allelic variants. As used herein, protein and polypeptide refers to a linear series of amino acid residues connected to one another by peptide bonds between the alpha-amino group and carboxy groups of adjacent amino

acid residues. Other covalent bonds, such as amide and disulfide bonds, may also be present.

The term "isolated" as used herein is considered to refer to molecules that are removed from their natural environment, i.e. isolated from a cell or from a living organism in which they normally occur, and that are separated or essentially purified from the coexisting components with which they are found to be associated in nature. This notion further means that the sequences encoding such molecules can be linked by the hand of man to polynucleotides, to which they are not linked in their natural state, and that such molecules can be produced by recombinant and/or synthetic means. Even if for said purposes those sequences may be introduced into living or non-living organisms by methods known to those skilled in the art, and even if those sequences are still present in said organisms, they are still considered to be isolated. In the present invention, the terms "risk", "susceptibility", and "predisposition" are tantamount and are used with respect to the probability of developing a neurodegenerative disease, preferably Alzheimer's disease.

The term 'AD' shall mean Alzheimer's disease. "AD-type neuropathology" as used herein refers to neuropathological, neurophysiological, histopathological and clinical hallmarks as described in the instant invention and as commonly known from state-of-the-art literature (see: Iqbal, Swaab, Winblad and Wisniewski, *Alzheimer's Disease and Related Disorders (Etiology, Pathogenesis and Therapeutics)*, Wiley & Sons, New York, Weinheim, Toronto, 1999; Scinto and Daffner, *Early Diagnosis of Alzheimer's Disease*, Humana Press, Totowa, New Jersey, 2000; Mayeux and Christen, *Epidemiology of Alzheimer's Disease: From Gene to Prevention*, Springer Press, Berlin, Heidelberg, New York, 1999; Younkin, Tanzi and Christen, *Presenilins and Alzheimer's Disease*, Springer Press, Berlin, Heidelberg, New York, 1998).

Neurodegenerative diseases or disorders according to the present invention comprise Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, Pick's disease, fronto-temporal dementia, progressive nuclear palsy, corticobasal degeneration, cerebro-vascular dementia, multiple system atrophy, argyrophilic grain dementia and other tauopathies, and mild-cognitive impairment. Further conditions involving neurodegenerative processes are, for instance, age-related macular degeneration, narcolepsy, motor neuron diseases, prion diseases, traumatic nerve injury and repair, and multiple sclerosis.

In one aspect, the invention features a method of diagnosing or prognosticating a neurodegenerative disease in a subject, or determining whether a subject is at increased risk of developing said disease. The method comprises: determining a level,

or an activity, or both said level and said activity of (i) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or of (ii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or of (iii) a fragment, or derivative, or variant of said transcription or translation product in a sample from said subject and comparing said level, and/or said activity to a reference value representing a known disease or health status, thereby diagnosing or prognosticating said neurodegenerative disease in said subject, or determining whether said subject is at increased risk of developing said neurodegenerative disease.

The invention also relates to the construction and the use of primers and probes which are unique to the nucleic acid sequences, or fragments, or variants thereof, as disclosed in the present invention. The oligonucleotide primers and/or probes can be labeled specifically with fluorescent, bioluminescent, magnetic, or radioactive substances. The invention further relates to the detection and the production of said nucleic acid sequences, or fragments and variants thereof, using said specific oligonucleotide primers in appropriate combinations. PCR-analysis, a method well known to those skilled in the art, can be performed with said primer combinations to amplify said gene specific nucleic acid sequences from a sample containing nucleic acids. Such sample may be derived either from healthy or diseased subjects. Whether an amplification results in a specific nucleic acid product or not, and whether a fragment of different length can be obtained or not, may be indicative for a neurodegenerative disease, in particular Alzheimer's disease. Thus, the invention provides nucleic acid sequences, oligonucleotide primers, and probes of at least 10 bases in length up to the entire coding and gene sequences, useful for the detection of gene mutations and single nucleotide polymorphisms in a given sample comprising nucleic acid sequences to be examined, which may be associated with neurodegenerative diseases, in particular Alzheimer's disease. This feature has utility for developing rapid DNA-based diagnostic tests, preferably also in the format of a kit.

In a further aspect, the invention features a method of monitoring the progression of a neurodegenerative disease in a subject. A level, or an activity, or both said level and said activity, of (i) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or of (ii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or of (iii) a fragment, or derivative, or variant of said transcription or translation product in a sample from said subject is determined. Said level and/or said activity is compared to a reference value representing a known disease or health status. Thereby the progression of said neurodegenerative disease in said subject is monitored.

In still a further aspect, the invention features a method of evaluating a treatment for a neurodegenerative disease, comprising determining a level, or an activity, or both said level and said activity of (i) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or of (ii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or of (iii) a fragment, or derivative, or variant of said transcription or translation product in a sample obtained from a subject being treated for said disease. Said level, or said activity, or both said level and said activity are compared to a reference value representing a known disease or health status, thereby evaluating the treatment for said neurodegenerative disease.

In a preferred embodiment of the herein claimed methods, kits, recombinant animals, molecules, assays, and uses of the instant invention, said gene coding for the membrane-associated guanylate kinase-interacting proteins is the gene coding for the human neuronal membrane-associated guanylate kinase-interacting protein 1 (SEQ ID NO. 5), also termed (MAGUIN-1), and the human neuronal membrane-associated guanylate kinase-interacting protein 2 (SEQ ID NO. 6).

In a further preferred embodiment of the herein claimed methods, kits, recombinant animals, molecules, assays, and uses of the instant invention, said neurodegenerative disease or disorder is Alzheimer's disease, and said subjects suffer from Alzheimer's disease.

The present invention discloses the differential expression and regulation of the human MAGUIN-1 and/or human MAGUIN-2 gene in specific brain regions of Alzheimer's disease patients. Consequently, human MAGUIN-1 and/or human MAGUIN-2 and their corresponding translation products may have a causative role in the regional selective neuronal degeneration typically observed in Alzheimer's disease. Alternatively, human MAGUIN-1 and/or human MAGUIN-2 may confer a neuroprotective function to the remaining surviving nerve cells. Based on these disclosures, the present invention has utility for the diagnostic evaluation and prognosis as well as for the identification of a predisposition to a neurodegenerative disease, in particular Alzheimer's disease. Furthermore, the present invention provides methods for the diagnostic monitoring of patients undergoing treatment for such a disease.

It is preferred that the sample to be analyzed and determined is selected from the group comprising brain tissue, or other tissues, or other body cells. The sample can also comprise cerebrospinal fluid or other body fluids including saliva, urine, serum plasma,

or mucus. Preferably, the methods of diagnosis, prognosis, monitoring the progression or evaluating a treatment for a neurodegenerative disease, according to the instant invention, can be practiced *ex corpore*, and such methods preferably relate to samples, for instance, body fluids or cells, removed, collected, or isolated from a subject or patient.

In further preferred embodiments, said reference value is that of a level, or an activity, or both said level and said activity of (i) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or of (ii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or of (iii) a fragment, or derivative, or variant of said transcription or translation product in a sample from a subject not suffering from said neurodegenerative disease.

In preferred embodiments, an alteration in the level and/or activity of a transcription product of the gene coding for human MAGUIN-1 and/or human MAGUIN-2 and/or a translation product of the gene coding for human MAGUIN-1 and/or human MAGUIN-2 protein in a sample cell, or tissue, or body fluid from said subject relative to a reference value representing a known health status indicates a diagnosis, or prognosis, or increased risk of becoming diseased with a neurodegenerative disease, particularly Alzheimer's disease.

In preferred embodiments, measurement of the level of transcription products of a gene coding for human MAGUIN-1 and/or human MAGUIN-2 is performed in a sample from a subject using a quantitative PCR-analysis with primer combinations to amplify said gene specific sequences from cDNA obtained by reverse transcription of RNA extracted from a sample of a subject. A Northern blot with probes specific for said gene can also be applied. It might also be preferred to measure transcription products by means of chip-based micro-array technologies. These techniques are known to those of ordinary skill in the art (see Sambrook and Russell, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2001; Schena M., *Microarray Biochip Technology*, Eaton Publishing, Natick, MA, 2000). An example of an immunoassay is the detection and measurement of enzyme activity as disclosed and described in the patent application WO 02/14543.

Furthermore, the level and/or an activity of a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or a fragment, or derivative, or variant of said translation product, and/or the level of activity of said translation product, and/or a fragment, or derivative, or variant thereof, can be detected using an immunoassay, an

activity assay, and/or a binding assay. These assays can measure the amount of binding between said protein molecule and an anti-protein antibody by the use of enzymatic, chromodynamic, radioactive, magnetic, or luminescent labels which are attached to either the anti-protein antibody or a secondary antibody which binds the anti-protein antibody. In addition, other high affinity ligands may be used. Immunoassays which can be used include e.g. ELISAs, Western blots and other techniques known to those of ordinary skill in the art (see Harlow and Lane, *Using Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1999 and Edwards R, *Immunodiagnosics: A Practical Approach*, Oxford University Press, Oxford; England, 1999). All these detection techniques may also be employed in the format of microarrays, protein-arrays, antibody microarrays, tissue microarrays, electronic biochip or protein-chip based technologies (see Schena M., *Microarray Biochip Technology*, Eaton Publishing, Natick, MA, 2000).

In a preferred embodiment, the level, or the activity, or both said level and said activity of (i) a transcription product of human MAGUIN-1 and/or human MAGUIN-2, and/or of (ii) a translation product of human MAGUIN-1 and/or human MAGUIN-2, and/or of (iii) a fragment, or derivative, or variant of said transcription or translation product in a series of samples taken from said subject over a period of time is compared, in order to monitor the progression of said disease. In further preferred embodiments, said subject receives a treatment prior to one or more of said sample gatherings. In yet another preferred embodiment, said level and/or activity is determined before and after said treatment of said subject.

In another aspect, the invention features a kit for diagnosing or prognosticating neurodegenerative diseases, in particular Alzheimer's disease, in a subject, or determining the propensity or predisposition of a subject to develop a neurodegenerative disease, in particular Alzheimer's disease, said kit comprising:

- (a) at least one reagent which is selected from the group consisting of (i) reagents that selectively detect a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2 (ii) reagents that selectively detect a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2; and
- (b) instruction for diagnosing, or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, or determining the propensity or predisposition of a subject to develop such a disease by
 - detecting a level, or an activity, or both said level and said activity, of said transcription product and/or said translation product of a gene coding for

human MAGUIN-1 and/or human MAGUIN-2, in a sample from said subject;
and

- diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, or determining the propensity or predisposition of said subject to develop such a disease,

wherein a varied level, or activity, or both said level and said activity, of said transcription product and/or said translation product compared to a reference value representing a known health status; or a level, or activity, or both said level and said activity, of said transcription product and/or said translation product similar or equal to a reference value representing a known disease status, indicates a diagnosis or prognosis of a neurodegenerative disease, in particular Alzheimer's disease, or an increased propensity or predisposition of developing such a disease. The kit, according to the present invention, may be particularly useful for the identification of individuals that are at risk of developing a neurodegenerative disease, in particular Alzheimer's disease. Consequently, the kit, according to the invention, may serve as a means for targeting identified individuals for early preventive measures or therapeutic intervention prior to disease onset, before irreversible damage in the course of the disease has been inflicted. Furthermore, in preferred embodiments, the kit featured in the invention is useful for monitoring a progression of a neurodegenerative disease, in particular Alzheimer's disease, in a subject, as well as monitoring success or failure of therapeutic treatment for such a disease of said subject.

In another aspect, the invention features a method of treating or preventing a neurodegenerative disease, in particular Alzheimer's disease, in a subject comprising the administration to said subject in a therapeutically or prophylactically effective amount of an agent or agents which directly or indirectly affect a level, or an activity, or both said level and said activity, of (i) a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (ii) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iii) a translation product of said gene, and/or (iv) a fragment, or derivative, or variant of (i) to (iii). Said agent may comprise a small molecule, or it may also comprise a peptide, an oligopeptide, or a polypeptide. Said peptide, oligopeptide, or polypeptide may comprise an amino acid sequence shown in SEQ ID NO.1 and/or SEQ ID NO.2, or a fragment, or derivative, or variant thereof. An agent for treating or preventing a neurodegenerative disease, in particular AD, according to the instant invention, may also consist of a nucleotide, an oligonucleotide, or a polynucleotide. Said oligonucleotide or polynucleotide may comprise a nucleotide sequence of the gene coding for human MAGUIN-1 and/or human MAGUIN-2 protein, either in sense orientation or in antisense orientation.

In preferred embodiments, the method comprises the application of per se known methods of gene therapy and/or antisense nucleic acid technology to administer said agent or agents. In general, gene therapy includes several approaches: molecular replacement of a mutated gene, addition of a new gene resulting in the synthesis of a therapeutic protein, and modulation of endogenous cellular gene expression by recombinant expression methods or by drugs. Gene-transfer techniques are described in detail (see e.g. Behr, *Acc Chem Res* 1993, 26: 274-278 and Mulligan, *Science* 1993, 260: 926-931; the contents of which are incorporated herein by reference) and include direct gene-transfer techniques such as mechanical microinjection of DNA into a cell as well as indirect techniques employing biological vectors (like recombinant viruses, especially retroviruses) or model liposomes, or techniques based on transfection with DNA coprecipitation with polycations, cell membrane perturbation by chemical (solvents, detergents, polymers, enzymes) or physical means (mechanic, osmotic, thermic, electric shocks). The postnatal gene transfer into the central nervous system has been described in detail (see e.g. Wolff, *Curr Opin Neurobiol* 1993, 3: 743-748).

In particular, the invention features a method of treating or preventing a neurodegenerative disease by means of antisense nucleic acid therapy, i.e. the down-regulation of an inappropriately expressed or defective gene by the introduction of antisense nucleic acids or derivatives thereof into certain critical cells (see e.g. Gillespie, *DN&P* 1992, 5: 389-395; Agrawal and Akhtar, *Trends Biotechnol* 1995, 13: 197-199; Crooke, *Biotechnology* 1992, 10: 882-6). Apart from hybridization strategies, the application of ribozymes, i.e. RNA molecules that act as enzymes, destroying RNA that carries the message of disease has also been described (see e.g. Barinaga, *Science* 1993, 262: 1512-1514). In preferred embodiments, the subject to be treated is a human, and therapeutic antisense nucleic acids or derivatives thereof are directed against human MAGUIN-1 and/or human MAGUIN-2. It is preferred that cells of the central nervous system, preferably the brain, of a subject are treated in such a way. Cell penetration can be performed by known strategies such as coupling of antisense nucleic acids and derivatives thereof to carrier particles, or the above described techniques. Strategies for administering targeted therapeutic oligodeoxynucleotides are known to those of skill in the art (see e.g. Wickstrom, *Trends Biotechnol* 1992, 10: 281-287). In some cases, delivery can be performed by mere topical application. Further approaches are directed to intracellular expression of antisense RNA. In this strategy, cells are transformed *ex vivo* with a recombinant gene that directs the synthesis of an RNA that is complementary to a region of target nucleic acid. Therapeutical use of intracellularly expressed antisense RNA is procedurally similar to gene therapy. A recently developed

method of regulating the intracellular expression of genes by the use of double-stranded RNA, known variously as RNA interference (RNAi), can be another effective approach for nucleic acid therapy (Hannon, *Nature* 2002, 418: 244-251).

In further preferred embodiments, the method comprises grafting donor cells into the central nervous system, preferably the brain, of said subject, or donor cells preferably treated so as to minimize or reduce graft rejection, wherein said donor cells are genetically modified by insertion of at least one transgene encoding said agent or agents. Said transgene might be carried by a viral vector, in particular a retroviral vector. The transgene can be inserted into the donor cells by a nonviral physical transfection of DNA encoding a transgene, in particular by microinjection. Insertion of the transgene can also be performed by electroporation, chemically mediated transfection, in particular calcium phosphate transfection, or liposomal mediated transfection (see Mc Celland and Pardee, *Expression Genetics: Accelerated and High-Throughput Methods*, Eaton Publishing, Natick, MA, 1999).

In preferred embodiments, said agent for treating and preventing a neurodegenerative disease, in particular AD, is a therapeutic protein which can be administered to said subject, preferably a human, by a process comprising introducing subject cells into said subject, said subject cells having been treated *in vitro* to insert a DNA segment encoding said therapeutic protein, said subject cells expressing *in vivo* in said subject a therapeutically effective amount of said therapeutic protein. Said DNA segment can be inserted into said cells *in vitro* by a viral vector, in particular a retroviral vector.

Methods of treatment, according to the present invention, comprise the application of therapeutic cloning, transplantation, and stem cell therapy using embryonic stem cells or embryonic germ cells and neuronal adult stem cells, combined with any of the previously described cell- and gene therapeutic methods. Stem cells may be totipotent or pluripotent. They may also be organ-specific. Strategies for repairing diseased and/or damaged brain cells or tissue comprise (i) taking donor cells from an adult tissue. Nuclei of those cells are transplanted into unfertilized egg cells from which the genetic material has been removed. Embryonic stem cells are isolated from the blastocyst stage of the cells which underwent somatic cell nuclear transfer. Use of differentiation factors then leads to a directed development of the stem cells to specialized cell types, preferably neuronal cells (Lanza et al., *Nature Medicine* 1999, 9: 975-977), or (ii) purifying adult stem cells, isolated from the central nervous system, or from bone marrow (mesenchymal stem cells), for *in vitro* expansion and subsequent grafting and transplantation, or (iii) directly inducing endogenous neural stem cells to proliferate,

migrate, and differentiate into functional neurons (Peterson DA, *Curr. Opin. Pharmacol.* 2002, 2: 34-42). Adult neural stem cells are of great potential for repairing damaged or diseased brain tissues, as the germinal centers of the adult brain are free of neuronal damage or dysfunction (Colman A, *Drug Discovery World* 2001, 7: 66-71).

In preferred embodiments, the subject for treatment or prevention, according to the present invention, can be a human, an experimental animal, e.g. a mouse or a rat, a domestic animal, or a non-human primate. The experimental animal can be an animal model for a neurodegenerative disorder, e.g. a transgenic mouse and/or a knock-out mouse with a neurodegenerative phenotype, in particular with an Alzheimer's-type neuropathology.

In a further aspect, the invention features a modulator of an activity, or a level, or both said activity and said level of at least one substance which is selected from the group consisting of (i) a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (ii) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2 and/or (iii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iv) a fragment, or derivative, or variant of (i) to (iii).

In an additional aspect, the invention features a pharmaceutical composition comprising said modulator and preferably a pharmaceutical carrier. Said carrier refers to a diluent, adjuvant, excipient, or vehicle with which the modulator is administered.

In a further aspect, the invention features a modulator of an activity, or a level, or both said activity and said level of at least one substance which is selected from the group consisting of (i) a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (ii) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iv) a fragment, or derivative, or variant of (i) to (iii) for use in a pharmaceutical composition.

In another aspect, the invention provides for the use of a modulator of an activity, or a level, or both said activity and said level of at least one substance which is selected from the group consisting of (i) a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (ii) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2 and/or (iii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iv) a fragment, or derivative, or variant of

(i) to (iii) for a preparation of a medicament for treating or preventing a neurodegenerative disease, in particular Alzheimer's disease.

In one aspect, the present invention also provides a kit comprising one or more containers filled with a therapeutically or prophylactically effective amount of said pharmaceutical composition.

In a further aspect, the invention features a recombinant, non-human animal comprising a non-native gene sequence coding for human MAGUIN-1 and/or human MAGUIN-2, or a fragment, or a variant, or a derivative thereof. The generation of said recombinant, non-human animal comprises (i) providing a gene targeting construct containing said gene sequence and a selectable marker sequence, and (ii) introducing said targeting construct into a stem cell of a non-human animal, and (iii) introducing said non-human animal stem cell into a non-human embryo, and (iv) transplanting said embryo into a pseudopregnant non-human animal, and (v) allowing said embryo to develop to term, and (vi) identifying a genetically altered non-human animal whose genome comprises a modification of said gene sequence in both alleles, and (vii) breeding the genetically altered non-human animal of step (vi) to obtain a genetically altered non-human animal whose genome comprises a modification of said endogenous gene, wherein said gene is mis-expressed, or under-expressed, or over-expressed, and wherein said disruption or alteration results in said non-human animal exhibiting a predisposition to developing symptoms of neuropathology similar to a neurodegenerative disease, in particular Alzheimer's disease. Strategies and techniques for the generation and construction of such an animal are known to those of ordinary skill in the art (see e.g. Capecchi, *Science* 1989, 244: 1288-1292 and Hogan et al., 1994, *Manipulating the Mouse Embryo: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York and Jackson and Abbott, *Mouse Genetics and Transgenics: A Practical Approach*, Oxford University Press, Oxford, England, 1999). It is preferred to make use of such a recombinant non-human animal as an animal model for investigating neurodegenerative diseases, in particular Alzheimer's disease. Such an animal may be useful for screening, testing and validating compounds, agents and modulators in the development of diagnostics and therapeutics to treat neurodegenerative diseases, in particular Alzheimer's disease.

In another aspect, the invention features an assay for screening for a modulator of neurodegenerative diseases, in particular Alzheimer's disease, or related diseases and disorders of one or more substances selected from the group consisting of (i) a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (ii) a transcription

product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iv) a fragment, or derivative, or variant of (i) to (iii). This screening method comprises (a) contacting a cell with a test compound, and (b) measuring the activity, or the level, or both the activity and the level of one or more substances recited in (i) to (iv), and (c) measuring the activity, or the level, or both the activity and the level of said substances in a control cell not contacted with said test compound, and (d) comparing the levels of the substance in the cells of step (b) and (c), wherein an alteration in the activity and/or level of said substances in the contacted cells indicates that the test compound is a modulator of said diseases and disorders.

In one further aspect, the invention features a screening assay for a modulator of neurodegenerative diseases, in particular Alzheimer's disease, or related diseases and disorders of one or more substances selected from the group consisting of (i) a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (ii) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iv) a fragment, or derivative, or variant of (i) to (iii), comprising (a) administering a test compound to a test animal which is predisposed to developing or has already developed symptoms of a neurodegenerative disease or related diseases or disorders, and (b) measuring the activity and/or level of one or more substances recited in (i) to (iv), and (c) measuring the activity and/or level of said substances in a matched control animal which is equally predisposed to developing or has already developed said symptoms and to which animal no such test compound has been administered, and (d) comparing the activity and/or level of the substance in the animals of step (b) and (c), wherein an alteration in the activity and/or level of substances in the test animal indicates that the test compound is a modulator of said diseases and disorders.

In a preferred embodiment, said test animal and/or said control animal is a recombinant, non-human animal which expresses the gene coding for human MAGUIN-1 and/or human MAGUIN-2, or a fragment, or derivative, or a variant thereof, under the control of a transcriptional regulatory element which is not the native human MAGUIN-1 and/or human MAGUIN-2 gene transcriptional control regulatory element.

In another embodiment, the present invention provides a method for producing a medicament comprising the steps of (i) identifying a modulator of neurodegenerative diseases by a method of the aforementioned screening assays and (ii) admixing the

modulator with a pharmaceutical carrier. However, said modulator may also be identifiable by other types of screening assays.

In another aspect, the present invention provides for an assay for testing a compound, preferably for screening a plurality of compounds, for inhibition of binding between a ligand and human MAGUIN-1 and/or human MAGUIN-2, or a fragment, or derivative, or variant thereof. Said screening assay comprises the steps of (i) adding a liquid suspension of said human MAGUIN-1 and/or human MAGUIN-2, or a fragment, or derivative, or variant thereof, to a plurality of containers, and (ii) adding a compound or a plurality of compounds to be screened for said inhibition to said plurality of containers, and (iii) adding fluorescently labelled ligand to said containers, and (iv) incubating said human MAGUIN-1 and/or human MAGUIN-2, or said fragment, or derivative, or variant thereof, and said compound or plurality of compounds, and said fluorescently labelled ligand, and (v) measuring the amounts of fluorescence associated with said human MAGUIN-1 and/or human MAGUIN-2, or with said fragment, or derivative, or variant thereof, and (vi) determining the degree of inhibition by one or more of said compounds of binding of said ligand to said human MAGUIN-1 and/or human MAGUIN-2, or said fragment, or derivative, or variant thereof. Instead of utilizing a fluorescently labelled ligand, it might in some aspects be preferred to use any other detectable label known to the person skilled in the art, e.g. radioactive labels, and detect it accordingly. Said method may be useful for the identification of novel compounds as well as for evaluating compounds which have been improved or otherwise optimized in their ability to inhibit the binding of a ligand to a gene product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, or a fragment, or derivative, or variant thereof. One example of a fluorescent binding assay, in this case based on the use of carrier particles, is disclosed and described in patent application WO 00/52451. A further example is the competitive assay method as described in patent WO 02/01226. Preferred signal detection methods for screening assays of the instant invention are described in the following patent applications: WO 96/13744, WO 98/16814, WO 98/23942, WO 99/17086, WO 99/34195, WO 00/66985, WO 01/59436, WO 01/59416.

In one further embodiment, the present invention provides a method for producing a medicament comprising the steps of (i) identifying a compound as an inhibitor of binding between a ligand and a gene product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2 by the aforementioned inhibitory binding assay and (ii) admixing the compound with a pharmaceutical carrier. However, said compound may also be identifiable by other types of screening assays.

In another aspect, the invention features an assay for testing a compound, preferably for screening a plurality of compounds to determine the degree of binding of said compounds to human MAGUIN-1 and/or human MAGUIN-2, or to a fragment, a variant, or derivative thereof. Said screening assay comprises (i) adding a liquid suspension of said human MAGUIN-1 and/or human MAGUIN-2, or a fragment, or a variant, or derivative thereof, to a plurality of containers, and (ii) adding a fluorescently labelled compound or a plurality of fluorescently labelled compounds to be screened for said binding to said plurality of containers, and (iii) incubating said human MAGUIN-1 and/or human MAGUIN-2, or said fragment, or variant, or derivative thereof, and said fluorescently labelled compound or fluorescently labelled compounds, and (iv) measuring the amounts of fluorescence associated with said human MAGUIN-1 and/or human MAGUIN-2, or with said fragment, or variant, or derivative thereof, and (v) determining the degree of binding by one or more of said compounds to said human MAGUIN-1 and/or human MAGUIN-2, or said fragment, or variant, or derivative thereof. In this type of assay it might be preferred to use a fluorescent label. However, any other type of detectable label might also be employed. Said method may be useful for the identification of novel compounds as well as for evaluating compounds which have been improved or otherwise optimized in their ability to bind to a human MAGUIN-1 and/or human MAGUIN-2 gene product, or fragment, or variant, or derivative thereof.

In one further embodiment, the present invention provides a method for producing a medicament comprising the steps of (i) identifying a compound as a binder to a gene product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2 by the aforementioned binding assays and (ii) admixing the compound with a pharmaceutical carrier. However, said compound may also be identifiable by other types of screening assays.

In another embodiment, the present invention provides for a medicament obtainable by any of the methods according to the herein claimed screening assays. In one further embodiment, the instant invention provides for a medicament obtained by any of the methods according to the herein claimed screening assays.

The present invention features protein molecules shown in SEQ ID NO. 1, and SEQ ID NO. 2, or fragments, or derivatives, or variants thereof, for use as diagnostic targets for detecting a neurodegenerative disease, preferably Alzheimer's disease.

The present invention further features protein molecules shown in SEQ ID NO. 1 and SEQ ID NO. 2, or fragments, or derivatives, or variants thereof, for use as screening

targets for reagents or compounds preventing, or treating, or ameliorating a neurodegenerative disease, preferably Alzheimer's disease.

The invention further features an antibody specifically immunoreactive with an immunogen, wherein said immunogen is a translation product of the human MAGUIN-1 gene shown in SEQ ID NO. 1, or a fragment, or a variant, or a derivative thereof. The immunogen may comprise immunogenic or antigenic epitopes or portions of a translation product of said gene, wherein said immunogenic or antigenic portion of a translation product is a polypeptide, and wherein said polypeptide elicits an antibody response in an animal, and wherein said polypeptide is immunospecifically bound by said antibody. Methods for generating antibodies are well known in the art (see Harlow et al., *Antibodies, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1988). The term "antibody", as employed in the present invention, encompasses all forms of antibodies known in the art, such as polyclonal, monoclonal, chimeric, recombinatorial, anti-idiotypic, humanized, or single chain antibodies, as well as fragments thereof (see Dubel and Breitling, *Recombinant Antibodies*, Wiley-Liss, New York, NY, 1999). Antibodies of the present invention are useful, for instance, in a variety of diagnostic and therapeutic methods, based on state-in-the-art techniques (see Harlow and Lane, *Using Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1999 and Edwards R., *Immunodiagnosics: A Practical Approach*, Oxford University Press, Oxford, England, 1999) such as enzyme-immuno assays (e.g. enzyme-linked immunosorbent assay, ELISA), radioimmuno assays, chemoluminescence-immuno assays, Western-blot, immunoprecipitation and antibody microarrays. These methods involve the detection of translation products of the human MAGUIN-1 gene.

The invention also features an antibody specifically immunoreactive with an immunogen, wherein said immunogen is a translation product of the human MAGUIN-2 gene shown in SEQ ID NO. 2, or a fragment, a variant, or a derivative thereof.

In a preferred embodiment of the present invention, said antibodies can be used for detecting the pathological state of a cell in a sample from a subject, comprising immunocytochemical staining of said cell with said antibody, wherein an altered degree of staining, or an altered staining pattern in said cell compared to a cell representing a known health status indicates a pathological state of said cell. The invention is particularly suited to detect pathological structures in the brain of a subject. It is also especially suited to detect pathological cells of the muscular system, prostate, stomach, testis, ovary, adrenal glands, mammary glands, liver, spleen, lung, trachea or placenta.

Preferably, the pathological state relates to a neurodegenerative disease, in particular to Alzheimer's disease. Immunocytochemical staining of a cell can be carried out by a number of different experimental methods well known in the art. It might be preferred, however, to apply an automated method for the detection of antibody binding, wherein the determination of the degree of staining of a cell, or the determination of the cellular or subcellular staining pattern of a cell, or the topological distribution of an antigen on the cell surface or among organelles and other subcellular structures within the cell, are carried out according to the method described in US patent 6150173.

Other features and advantages of the invention will be apparent from the following description of figures and examples which are illustrative only and not intended to limit the remainder of the disclosure in any way.

Figure 1 depicts the brain regions with selective vulnerability to neuronal loss and degeneration in Alzheimer's disease. Primarily, neurons within the inferior temporal lobe, the entorhinal cortex, the hippocampus, and the amygdala are subject to degenerative processes in Alzheimer's disease (Terry et al., *Annals of Neurology* 1981, 10:184-192). These brain regions are mostly involved in the processing of learning and memory functions. In contrast, neurons within the frontal cortex, the occipital cortex, and the cerebellum remain largely intact and preserved from neurodegenerative processes in Alzheimer's disease. Brain tissues from the frontal cortex (F), the temporal cortex (T), and the hippocampus (H) of Alzheimer's disease patients and healthy, age-matched control individuals were used for the herein disclosed examples. For illustrative purposes, the image of a normal healthy brain was taken from a publication by Strange (*Brain Biochemistry and Brain Disorders*, Oxford University Press, Oxford, 1992, p.4).

Figure 2 discloses the initial identification of the differential expression of the human MAGUIN-1 gene in a fluorescence differential display screen. The figure shows a clipping of a large preparative fluorescent differential display gel. PCR products from the frontal cortex (F) and the temporal cortex (T) of two healthy control subjects and six Alzheimer's diseased patients were loaded in duplicate onto a denaturing polyacrylamide gel (from left to right). PCR products were obtained by amplification of the individual cDNAs with the corresponding one-base-anchor oligonucleotide and the specific Cy3 labelled random primers. The arrow indicates the lane where significant differences in intensity of the signals for human MAGUIN-1 transcript derived from frontal cortex, compared to the signals for human MAGUIN-1 transcript derived from the temporal cortex of Alzheimer's patients exist. The differential expression reflects a down-regulation of human MAGUIN RNA expression in the temporal cortex in

comparison to the frontal cortex of AD patients. Comparing the signals derived from frontal cortex and temporal cortex of healthy non-AD control subjects with each other, no distinction in signal intensity, i.e. no altered expression level can be detected.

Figure 3 and Figure 4 illustrate the verification of the differential expression of human MAGUIN-1 in AD brain tissues by quantitative RT-PCR analysis. Quantification of RT-PCR products from RNA samples collected from the frontal cortex (F) and the temporal cortex (T) of AD patients (Figure 3a) and samples from the frontal cortex (F) and the hippocampus (H) of AD patients (Figure 4a) was performed by the LightCycler rapid thermal cycling technique. Likewise, samples of healthy, age-matched control individuals were compared (Figure 3b for frontal cortex and temporal cortex, Figure 4b for frontal cortex and hippocampus). The data were normalized to the combined average values of a set of standard genes which showed no significant differences in their gene expression levels. Said set of standard genes consisted of genes for cyclophilin B, the ribosomal protein S9, the transferrin receptor, GAPDH, and beta-actin. The figure depicts the kinetics of amplification by plotting the cycle number against the amount of amplified material as measured by its fluorescence. Note that the amplification kinetics of Maguin-1 cDNA from both, the frontal and temporal cortices of a normal control individual, and from the frontal cortex and hippocampus of a normal control individual, respectively, during the exponential phase of the reaction are juxtaposed (Figures 3b and 4b, arrowheads), whereas in Alzheimer's disease (Figures 3a and 4a, arrowheads) there is a significant separation of the corresponding curves, indicating a differential expression of Maguin-1 in the respective analyzed brain regions.

Figure 5 and Figure 6 illustrate the verification of the differential expression of human MAGUIN-2 in AD brain tissues by quantitative RT-PCR analysis. Quantification of RT-PCR products from RNA samples collected from the frontal cortex (F) and the temporal cortex (T) of AD patients (Figure 5a) and samples from the frontal cortex (F) and the hippocampus (H) of AD patients (Figure 6a) was performed by the LightCycler rapid thermal cycling technique. Likewise, samples of healthy, age-matched control individuals were compared (Figure 5b for frontal cortex and temporal cortex, Figure 6b for frontal cortex and hippocampus). The data were normalized to the combined average values of a set of standard genes which showed no significant differences in their gene expression levels. Said set of standard genes consisted of genes for cyclophilin B, the ribosomal protein S9, the transferrin receptor, GAPDH, and beta-actin. The figure depicts the kinetics of amplification by plotting the cycle number against the amount of amplified material as measured by its fluorescence. Note that the amplification kinetics of Maguin-2 cDNA from both, the frontal and temporal cortices of a normal control

individual, and from the frontal cortex and hippocampus of a normal control individual, respectively, during the exponential phase of the reaction are juxtaposed (Figures 5b and 6b, arrowheads), whereas in Alzheimer's disease (Figures 5a and 6a, arrowheads) there is a significant separation of the corresponding curves, indicating a differential expression of Maguin-2 in the respective analyzed brain regions.

Figure 7 discloses the amino acid sequence of the human MAGUIN-1 protein comprising 1034 amino acids; SEQ ID NO. 1. The protein harbors several distinct functional domains which are located as follows: amino acid residues 8 to 75 form the 'Sterile Alpha Motif' (SAM), amino acid residues 156 to 296 constitute the PDZ domain, and the Pleckstrin-homology (PH) domain consists of amino acid residues 572 to 667.

Figure 8 shows an alignment of the amino acid sequence of SEQ ID NO.1, human MAGUIN-1, with the rat MAGUIN-1 amino acid sequence (GenBank accession number aad04568). The full length human MAGUIN-1 protein consists of 1034 amino acids (residues given in the single-letter amino acid code).

Figure 9 discloses the amino acid sequence of the human MAGUIN-2 protein comprising 898 amino acids; SEQ ID NO. 2. The protein harbors several distinct functional domains which are located as follows: amino acid residues 8 to 75 form the 'Sterile Alpha Motif' (SAM), amino acid residues 156 to 296 constitute the PDZ domain, and the Pleckstrin-homology (PH) domain consists of amino acid residues 572 to 667.

Figure 10 shows an alignment of the amino acid sequence of SEQ ID NO. 2, human MAGUIN-2, with rat MAGUIN-2 amino acid sequence (GenBank accession number aad04567). The full length human MAGUIN-2 protein consists of 898 amino acids (residues given in the single-letter amino acid code).

Figure 11 represents the nucleotide sequence of SEQ ID NO. 3, the coding sequence of the human MAGUIN-1 gene, comprising 3105 nucleotides.

Figure 12 represents the nucleotide sequence of SEQ ID NO. 4, the coding sequence of the human MAGUIN-2 gene, comprising 2697 nucleotides.

Figure 13 shows SEQ ID NO. 5, the nucleotide sequence of the human MAGUIN-1 cDNA, comprising 5749 nucleotides.

Figure 14 shows SEQ ID NO. 6, the nucleotide sequence of the human MAGUIN-2 cDNA; comprising 4350 nucleotides.

Figure 15 depicts SEQ ID NO. 7, the nucleotide sequence of the 50 bp MAGUIN-1 cDNA fragment, identified and obtained by fluorescence differential display and subsequent cloning.

Figure 16 outlines the sequence alignment of SEQ ID NO. 7, the 50 bp human MAGUIN-1 cDNA fragment, with the 3'UTR nucleotide sequence of SEQ ID NO. 5 (nucleotide 5693 to 5742), the nucleotide sequence of human MAGUIN-1 cDNA.

Figure 17 charts the schematic alignment of SEQ ID NO. 7, the human MAGUIN-1 cDNA fragment, SEQ ID NO. 6, the human MAGUIN-2 cDNA sequence and the nucleotide sequence of SEQ ID NO. 5, the nucleotide sequence of human MAGUIN-1 cDNA, derived from the alignment of EST nucleotide sequences as found in the GenBank genetic sequence database. EST numbers are written on the left side, all sequences are 5' to 3' directed.

Figure 18 depicts human cerebral cortex labeled with an affinity-purified rabbit anti-Maguin-1 antiserum raised against a peptide corresponding to amino acids 914-928 (green signals). Strong immunoreactivity of human Maguin-1 was detected in neuronal cell bodies (indicated by arrowheads) and in neurites, whereas glial cells were immunonegative (see arrows). The same immunostaining pattern was observed by using another antiserum raised against a peptide mapping to amino acids 973-987 of Maguin-1. Blue signals indicate nuclei stained with DAPI.

Table 1 lists the gene expression levels in the frontal cortex relative to the temporal cortex for the human MAGUIN-1 gene in seven Alzheimer's disease patients, herein identified by internal reference numbers P010, P011, P012, P014, P016, P017, P019 (1.42 to 4.14 fold) and five healthy, age-matched control individuals, herein identified by internal reference numbers C005, C008, C011, C012, C014 (0.30 to 1.36 fold). The values shown are reciprocal values according to the formula described herein.

Table 2 lists the gene expression levels in the frontal cortex relative to the hippocampus for the human MAGUIN-1 gene in six Alzheimer's disease patients, herein identified by internal reference numbers P010, P011, P012, P014, P016, P019 (1.21 to 3.07 fold) and three healthy, age-matched control individuals, herein identified by internal reference

numbers C004, C005, C008 (0.39 to 1.74 fold). The values shown are reciprocal values according to the formula described herein.

Table 3 lists the gene expression levels in the frontal cortex relative to the temporal cortex for the human MAGUIN-2 gene in seven Alzheimer's disease patients, herein identified by internal reference numbers P010, P011, P012, P014, P016, P017, P019 (1.77 to 11.73 fold) and five healthy, age-matched control individuals, herein identified by internal reference numbers C005, C008, C011, C012, C014 (0.30 to 1.42 fold). The values shown are reciprocal values according to the formula described herein.

Table 4 lists the gene expression levels in the frontal cortex relative to the hippocampus for the human MAGUIN-2 gene in six Alzheimer's disease patients, herein identified by internal reference numbers P010, P011, P012, P014, P016, P019 (0.72 to 9.08 fold) and three healthy, age-matched control individuals, herein identified by internal reference numbers C004, C005, C008 (0.46 to 1.69 fold). The values shown are reciprocal values according to the formula described herein.

EXAMPLE I:

(i) Brain tissue dissection from patients with Alzheimer's disease:

Brain tissues from Alzheimer's disease patients and age-matched control subjects were collected within 6 hours post-mortem and immediately frozen on dry ice. Sample sections from each tissue were fixed in paraformaldehyde for histopathological confirmation of the diagnosis. Brain areas for differential expression analysis were identified (see Figure 1) and stored at - 80 °C until RNA extractions were performed.

(ii) Isolation of total RNA:

Total RNA was extracted from post-mortem brain tissue by using the RNeasy kit (Qiagen) according to the manufacturer's protocol. The accurate RNA concentration and the RNA quality were determined with the DNA LabChip system using the Agilent 2100 Bioanalyzer (Agilent Technologies). For additional quality testing of the prepared RNA, i.e. exclusion of partial degradation and testing for DNA contamination, specifically designed intronic GAPDH oligonucleotides and genomic DNA as reference control were utilised to generate a melting curve with the LightCycler technology as described in the manufacturer's protocol (Roche).

(iii) cDNA synthesis and identification of differentially expressed genes by

fluorescence differential display (FDD):

In order to identify changes in gene expression in different tissues we employed a modified and improved differential display (DD) screening method. The original DD screening method is known to those skilled in the art (Liang and Pardee, *Science* 1995, 267:1186-7). This technique compares two populations of RNA and provides clones of genes that are expressed in one population but not in the other. Several samples can be analyzed simultaneously and both up- and down-regulated genes can be identified in the same experiment. By adjusting and refining several steps in the DD method as well as modifying technical parameters, e.g. increasing redundancy, evaluating optimized reagents and conditions for reverse transcription of total RNA, optimizing polymerase chain reactions (PCR) and separation of the products thereof, a technique was developed which allows for highly reproducible and sensitive results. The applied and improved DD technique was described in detail by von der Kammer et al. (*Nucleic Acids Research* 1999, 27: 2211-2218). A set of 64 specifically designed random primers were developed (standard set) to achieve a statistically comprehensive analysis of all possible RNA species. Further, the method was modified to generate a preparative DD slab-gel technique, based on the use of fluorescently labelled primers. In the present invention, RNA populations from carefully selected post-mortem brain tissues (frontal and temporal cortex) of Alzheimer's disease patients and age-matched control subjects were compared.

As starting material for the DD analysis we used total RNA, extracted as described above (ii). Equal amounts of 0.05 µg RNA each were transcribed into cDNA in 20 µl reactions containing 0.5 mM each dNTP, 1 µl SensiscriptTM Reverse Transcriptase and 1x RT buffer (Qiagen), 10 U RNase inhibitor (Qiagen) and 1 µM of either one-base-anchor oligonucleotides HT₁₁A, HT₁₁G or HT₁₁C (Liang et al., *Nucleic Acids Research* 1994, 22: 5763-5764; Zhao et al., *Biotechniques* 1995, 18: 842-850). Reverse transcription was performed for 60 min at 37 °C with a final denaturation step at 93 °C for 5 min. 2 µl of the obtained cDNA each was subjected to a polymerase chain reaction (PCR) employing the corresponding one-base-anchor oligonucleotide (1 µM) along with either one of the Cy3 labelled random DD primers (1 µM), 1x GeneAmp PCR buffer (Applied Biosystems), 1.5 mM MgCl₂ (Applied Biosystems), 2 µM dNTP-Mix (dATP, dGTP, dCTP, dTTP Amersham Pharmacia Biotech), 5 % DMSO (Sigma), 1 U AmpliTaq DNA Polymerase (Applied Biosystems) in a 20 µl final volume. PCR conditions were set as follows: one round at 94 °C for 30 sec for denaturing, cooling 1 °C/sec down to 40 °C, 40 °C for 4 min for low-stringency annealing of primer, heating 1 °C/sec up to 72 °C, 72 °C for 1 min for extension. This round was followed by 39 high-stringency cycles: 94

°C for 30 sec, cooling 1 °C/sec down to 60 °C, 60 °C for 2 min, heating 1 °C/sec up to 72 °C, 72 °C for 1 min. One final step at 72 °C for 5 min was added to the last cycle (PCR cycler: Multi Cycler PTC 200, MJ Research). 8 µl DNA loading buffer were added to the 20 µl PCR product preparation, denatured for 5 min and kept on ice until loading onto a gel. 3.5 µl each were separated on 0.4 mm thick, 6 %-polyacrylamide (Long Ranger)/ 7 M urea sequencing gels in a slab-gel system (Hitachi Genetic Systems) at 2000 V, 60W, 30 mA, for 1 h 40 min. Following completion of the electrophoresis, gels were scanned with a FMBIO II fluorescence-scanner (Hitachi Genetic Systems), using the appropriate FMBIO II Analysis 8.0 software. A full-scale picture was printed, differentially expressed bands marked, excised from the gel, transferred into 1.5 ml containers, overlayed with 200 µl sterile water and kept at -20°C until extraction.

Elution and reamplification of differential display products: The differential bands were extracted from the gel by boiling in 200 µl H₂O for 10 min, cooling down on ice and precipitation from the supernatant fluids by using ethanol (Merck) and glycogen/sodium acetate (Merck) at - 20 °C over night, and subsequent centrifugation at 13.000 rpm for 25 min at 4 °C. Pellets were washed twice in ice-cold ethanol (80%), resuspended in 10 mM Tris pH 8.3 (Merck) and dialysed against 10 % glycerol (Merck) for 1 h at room temperature on a 0.025 µm VSWP membrane (Millipore). The obtained preparations were used as templates for reamplification by 15 high-stringency cycles in 25-µl PCR mixtures containing the corresponding primer pairs as used for the differential display PCR (see above) under identical conditions, with the exception of the initial round at 94 °C for 5 min, followed by 15 cycles of: 94 °C for 45 sec, 60 °C for 45 sec, ramp 1°C/sec to 70 °C for 45 sec, and one final step at 72 °C for 5 min.

Cloning and sequencing of differential display products: Re-amplified cDNAs were analyzed with the DNA LabChip system (Agilent 2100 Bioanalyzer, Agilent Technologies) and were ligated into the pCR-Blunt II-TOPO vector and transformed into *E.coli* Top10F' cells (Zero Blunt TOPO PCR Cloning Kit, Invitrogen) according to the manufacturer's instructions. Cloned cDNA fragments were sequenced by commercially available sequencing facilities. The results of one such fluorescence differential display experiment for the human MAGUIN-1 gene are shown in Figure 2.

(iv) Confirmation of differential expression by quantitative RT-PCR:

Positive corroboration of differential expression of the human MAGUIN-1 gene and human MAGUIN-2 gene was performed using the LightCycler technology (Roche). This technique features rapid thermal cycling for the polymerase chain reaction as well as real-time measurement of fluorescent signals during amplification and therefore allows for highly accurate quantification of RT-PCR products by using a kinetic, rather than an endpoint approach. The ratios of human MAGUIN-1 and human MAGUIN-2 cDNA each

from the temporal cortex and frontal cortex, and from the hippocampus and frontal cortex, respectively, were determined (relative quantification).

First, a standard curve was generated to determine the efficiency of the PCR with specific primers for human MAGUIN-1:

5'-CAGCAAGCAGTTGACGGGA-3'

and 5'-GCCACGAGTCTTGTCAAATTCA -3'

and human MAGUIN-2:

5'-GGGCCTCCCAAAGGGATAT-3'

and 5'-CCCAATGTAGAAAGCTCGCATT-3'.

PCR amplification (95 °C and 1 sec, 56 °C and 5 sec, and 72 °C and 5 sec) was performed in a volume of 20 µl containing Lightcycler-FastStart DNA Master SYBR Green I mix (contains FastStart Taq DNA polymerase, reaction buffer, dNTP mix with dUTP instead of dTTP, SYBR Green I dye, and 1 mM MgCl₂; Roche), 0.5 µM primers, 2 µl of a cDNA dilution series (final concentration of 40, 20, 10, 5, 1 and 0.5 ng human total brain cDNA; Clontech) and depending on the primers used, additional 3 mM MgCl₂. Melting curve analysis revealed a single peak at approximately 80°C and 80.6°C with no visible primer dimers. Quality and size of the PCR product were determined with the DNA LabChip system (Agilent 2100 Bioanalyzer, Agilent Technologies). A single peak at the expected size of 121 bp for human MAGUIN-1 and 67 bp for human MAGUIN-2 was observed in the electropherogram of the sample.

In an analogous manner, the PCR protocol was applied to determine the PCR efficiency of a set of reference genes which were selected as a reference standard for quantification. In the present invention, the mean value of five such reference genes was determined: (1) cyclophilin B, using the specific primers 5'-ACTGAAGCACTACGGGCCTG-3' and 5'-AGCCGTTGGTGTCTTTGCC-3' except for MgCl₂ (an additional 1 mM was added instead of 3 mM). Melting curve analysis revealed a single peak at approximately 87 °C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band of the expected size (62 bp). (2) Ribosomal protein S9 (RPS9), using the specific primers 5'-GGTCAAATTTACCCTGGCCA-3' and 5'-TCTCATCAAGCGTCAGCAGTTC-3' (exception: additional 1 mM MgCl₂ was added instead of 3 mM). Melting curve analysis revealed a single peak at approximately 85°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (62 bp). (3) beta-actin, using the specific primers 5'-TGGAACGGTGAAGGTGACA-3' and 5'-GGCAAGGGACTTCCTGTAA-3'. Melting curve analysis revealed a single peak at approximately 87°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (142 bp). (4) GAPDH, using the specific primers 5'-CGTCATGGGTGTGAACCATG-

3' and 5'-GCTAAGCAGTTGGTGGTGCAG-3'. Melting curve analysis revealed a single peak at approximately 83°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (81 bp). (5) Transferrin receptor TRR, using the specific primers 5'-GTCGCTGGTCAGTTCGTGATT-3' and 5'-AGCAGTTGGCTGTTGTACCTCTC-3'. Melting curve analysis revealed a single peak at approximately 83°C with no visible primer dimers. Agarose gel analysis of the PCR product showed one single band with the expected size (80 bp).

For calculation of the values, first the logarithm of the cDNA concentration was plotted against the threshold cycle number C_t for human MAGUIN-1 and human MAGUIN-2, respectively, and the five reference standard genes. The slopes and the intercepts of the standard curves (i.e. linear regressions) were calculated for all genes. In a second step, cDNAs from temporal cortex and frontal cortex, and from hippocampus and frontal cortex, for human MAGUIN-1 and human MAGUIN-2, respectively, were analyzed in parallel and normalized to cyclophilin B. The C_t values were measured and converted to ng total brain cDNA using the corresponding standard curves:

$$10^{(C_t \text{ value} - \text{intercept}) / \text{slope}} \quad [\text{ng total brain cDNA}]$$

The values for temporal cortex and frontal cortex of human MAGUIN-1 and human MAGUIN-2 cDNAs, and the values for hippocampus and frontal cortex of human MAGUIN-1 and human MAGUIN-2 cDNAs, respectively, were normalized to cyclophilin B, and the ratios were calculated using the following formulas:

$$\text{Ratio} = \frac{\text{human MAGUIN-1(MAGUIN-2) temporal [ng]} / \text{cyclophilin B temporal [ng]}}{\text{human MAGUIN-1(MAGUIN-2) frontal [ng]} / \text{cyclophilin B frontal [ng]}}$$

$$\text{Ratio} = \frac{\text{human MAGUIN-1(MAGUIN-2) hippocampus [ng]} / \text{cyclophilin B hippocampus [ng]}}{\text{human MAGUIN-1(MAGUIN-2) frontal [ng]} / \text{cyclophilin B frontal [ng]}}$$

In a third step, the set of reference standard genes was analyzed in parallel to determine the mean average value of the temporal to frontal ratios, and of the hippocampal to frontal ratios, respectively, of expression levels of the reference standard genes for each individual brain sample. As cyclophilin B was analyzed in step

2 and step 3, and the ratio from one gene to another gene remained constant in different runs, it was possible to normalize the values for human MAGUIN-1 and human MAGUIN-2 to the mean average value of the set of reference standard genes instead of normalizing to one single gene alone. The calculation was performed by dividing the ratio shown above by the deviation of cyclophilin B from the mean value of all housekeeping genes. The results of such quantitative RT-PCR analysis for the human MAGUIN-1 gene are shown in Figure 3 and 4 and for the human MAGUIN-2 gene in Figure 5 and 6.

(v) Sequence Analysis

Searching the EST database of the GenBank database for sequence similarities to the identified differentially expressed human cDNA fragment, SEQ ID NO. 7, as stated in the present invention, it was found that SEQ ID NO. 7 was identical to portions of the human EST sequences ai817268 and bf115709, (shown in Figure 17). These human ESTs showed high homology to rat (*Rattus norvegicus*) MAGUIN-1. Aligning human ESTs in addition to the identified expressed SEQ ID NO. 7, a complete EST cluster representing the human MAGUIN-1 cDNA, SEQ ID NO. 5, was constructed. The amino acid sequence of a large open reading frame with the potential to encode a protein of 1034 amino acid residues was deduced (SEQ ID NO. 1). The human MAGUIN-1 protein, as denoted herein, is highly homologous to the rat MAGUIN-1 protein. In addition, it encodes a protein (SEQ ID NO. 1) harboring a number of structurally and functionally important domains. One SAM, one PDZ, and one PH domain are located from amino acid residues 8 to 75 (SAM), amino acid residues 156 to 296 (PDZ), and amino acid residues 572 to 667 (PH) (refer to Figure 7).

(vi) Immunohistochemistry:

For immunofluorescence staining of Maguin-1 in human brain, frozen sections were prepared from post-mortem pre-central gyrus of a donor person (Cryostat Leica CM3050S) and fixed in acetone for 10 min. After washing in PBS, the sections were pre-incubated with blocking buffer (10% normal goat serum, 0.2% Triton X-100 in PBS) for 30min, and then incubated with affinity-purified rabbit anti-Maguin-1 antisera (1:20-1:50 diluted in blocking buffer, custom-made by Davids Biotechnologie, Regensburg) overnight at 4°C. After rinsing three times in 0.2% Triton X-100/PBS, the sections were incubated with FITC-conjugated goat anti-rabbit IgG (1:150 diluted in 1% BSA/PBS) for 2 hours at room temperature, and then again washed in PBS. Staining of the nuclei was performed by incubation of the sections with 5µM DAPI in PBS for 3min (blue signal). In order to block the autofluorescence of lipofuscin in human brain, the sections were treated with 1% Sudan Black B in 70% ethanol for 2-10 min at room temperature,

sequentially dipped in 70% ethanol, distilled water and PBS. The sections were coverslipped by 'Vectrashield mounting medium' (Vector Laboratories, Burlingame, CA) and observed under an inverted microscope (IX81, Olympus Optical). The digital images were captured with the appropriate software (AnalySiS, Olympus Optical).

CLAIMS

1. A method of diagnosing or prognosticating a neurodegenerative disease in a subject, or determining whether a subject is at increased risk of developing said disease, comprising:

determining a level and/or an activity of

- (i) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or
- (ii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or
- (iii) a fragment, or derivative, or variant of said transcription or translation product,

in a sample from said subject and comparing said level and/or said activity to a reference value representing a known disease or health status, thereby diagnosing or prognosticating said neurodegenerative disease in said subject, or determining whether said subject is at increased risk of developing said neurodegenerative disease.

2. A method of monitoring the progression of a neurodegenerative disease in a subject, comprising:

determining a level and/or an activity of

- (i) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or
- (ii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or
- (iii) a fragment, or derivative, or variant of said transcription or translation product,

in a sample from said subject and comparing said level and/or said activity to a reference value representing a known disease or health status, thereby monitoring the progression of said neurodegenerative disease in said subject.

3. A method of evaluating a treatment for a neurodegenerative disease, comprising:

determining a level and/or an activity of

- (i) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or
- (ii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or

- (iii) a fragment, or derivative, or variant of said transcription or translation product,

in a sample from a subject being treated for said disease and comparing said level and/or said activity to a reference value representing a known disease or health status, thereby evaluating said treatment for said neurodegenerative disease.

4. The method according to any of claims 1 to 3 wherein said neurodegenerative disease is Alzheimer's disease.

5. The method according to any of claims 1 to 4 wherein said sample is a cell, or a tissue, or a body fluid, in particular cerebrospinal fluid or blood.

6. The method according to any of claims 1 to 5 wherein said reference value is that of a level and/or an activity of

- (i) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or

- (ii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or

- (iii) a fragment, or derivative, or variant of said transcription or translation product,

in a sample from a subject not suffering from said neurodegenerative disease.

7. The method according to any of claims 1 to 6 wherein an alteration in the level and/or activity of a transcription product of the gene coding for human MAGUIN-1 and/or human MAGUIN-2 and/or a translation product of the gene coding for human MAGUIN-1 and/or human MAGUIN-2 and/or a fragment, or derivative, or variant thereof, in a sample cell, or tissue, or body fluid, in particular cerebrospinal fluid, from said subject relative to a reference value representing a known health status indicates a diagnosis, or prognosis, or increased risk of Alzheimer's disease in said subject.

8. The method according to any of claims 1 to 7, further comprising comparing a level and/or an activity of

- (i) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or

- (ii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or

- (iii) a fragment, or derivative, or variant of said transcription or translation product,

in a series of samples taken from said subject over a period of time.

9. The method according to claim 8, wherein said subject receives a treatment prior to one or more of said sample gatherings.

10. The method according to claim 9 wherein said level and/or activity is determined before and after said treatment of said subject.

11. A kit for diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, in a subject, or determining the propensity or predisposition of a subject to develop such a disease, said kit comprising:

- (a) at least one reagent which is selected from the group consisting of
 - (i) reagents that selectively detect a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and
 - (ii) reagents that selectively detect a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2 and
- (b) an instruction for diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, or determining the propensity or predisposition of a subject to develop such a disease by
 - (i) detecting a level, or an activity, or both said level and said activity, of said transcription product and/or said translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2 in a sample from said subject; and
 - (ii) diagnosing or prognosticating a neurodegenerative disease, in particular Alzheimer's disease, or determining the propensity or predisposition of said subject to develop such a disease, wherein a varied level, or activity, or both said level and said activity, of said transcription product and/or said translation product compared to a reference value representing a known health status; or a level, or activity, or both said level and said activity, of said transcription product and/or said translation product similar or equal to a reference value representing a known disease status indicates a diagnosis or prognosis of a neurodegenerative disease, in particular Alzheimer's disease, or an increased propensity or predisposition of developing such a disease.

12. A method of treating or preventing a neurodegenerative disease, in particular Alzheimer's disease, in a subject comprising administering to said subject in a therapeutically or prophylactically effective amount an agent or agents which directly or

indirectly affect an activity and/or a level of (i) a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (ii) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iv) a fragment, or derivative, or variant of (i) to (iii).

13. A modulator of an activity and/or of a level of at least one substance which is selected from the group consisting of (i) a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (ii) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iv) a fragment, or derivative, or variant of (i) to (iii).

14. A pharmaceutical composition comprising a modulator according to claim 13.

15. Use of a modulator of an activity and/or of a level of at least one substance which is selected from the group consisting of (i) a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (ii) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or (iv) a fragment, or derivative, or variant of (i) to (iii) for a preparation of a medicament for treating or preventing a neurodegenerative disease, in particular Alzheimer's disease.

16. A recombinant, non-human animal comprising a non-native gene sequence coding for human MAGUIN-1 and/or human MAGUIN-2 or a fragment, or a derivative, or a variant thereof, said animal being obtainable by:

- (i) providing a gene targeting construct comprising said gene sequence and a selectable marker sequence, and
- (ii) introducing said targeting construct into a stem cell of a non-human animal, and
- (iii) introducing said non-human animal stem cell into a non-human embryo, and
- (iv) transplanting said embryo into a pseudopregnant non-human animal, and
- (v) allowing said embryo to develop to term, and
- (vi) identifying a genetically altered non-human animal whose genome comprises a modification of said gene sequence in both alleles, and
- (vii) breeding the genetically altered non-human animal of step (vi) to obtain a genetically altered non-human animal whose genome comprises a modification of said endogenous gene, wherein said disruption results in said

non-human animal exhibiting a predisposition to developing symptoms of a neurodegenerative disease or related diseases or disorders.

17. Use of the recombinant, non-human animal according to claim 16 for screening, testing, and validating compounds, agents, and modulators in the development of diagnostics and therapeutics to treat neurodegenerative diseases, in particular Alzheimer's disease.

18. An assay for screening for a modulator of neurodegenerative diseases, in particular Alzheimer's disease, or related diseases or disorders of one or more substances selected from the group consisting of

- (i) a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or
- (ii) a transcription product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or
- (iii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2, and/or
- (iv) a fragment, or derivative, or variant of (i) to (iii), said method comprising:
 - (a) contacting a cell with a test compound;
 - (b) measuring the activity and/or level of one or more substances recited in (i) to (iv);
 - (c) measuring the activity and/or level of one or more substances recited in (i) to (iv) in a control cell not contacted with said test compound; and
 - (d) comparing the levels and/or activities of the substance in the cells of step (b) and (c), wherein an alteration in the activity and/or level of substances in the contacted cells indicates that the test compound is a modulator of said diseases or disorders.

19. A method of testing a compound, preferably of screening a plurality of compounds, for inhibition of binding between a ligand and human MAGUIN-1 and/or human MAGUIN-2, or fragments, or derivatives, or variants thereof, said method comprising the steps of:

- (i) adding a liquid suspension of said human MAGUIN-1 and/or human MAGUIN-2, or fragments, or derivatives, or variants thereof, to a plurality of containers;
- (ii) adding a compound, preferably a plurality of compounds, to be screened for said inhibition of binding to said plurality of containers;
- (iii) adding a detectable ligand, in particular a fluorescently detectable ligand, to said containers;

- (iv) incubating the liquid suspension of said human MAGUIN-1 and/or human MAGUIN-2, or said fragments, or derivatives, or variants thereof, and said compound, preferably said plurality of compounds, and said ligand;
- (v) measuring amounts of detectable ligand or fluorescence associated with said human MAGUIN-1 and/or human MAGUIN-2, or with said fragments, or derivatives, or variants thereof; and
- (vi) determining the degree of inhibition by one or more of said compounds of binding of said ligand to said human MAGUIN-1 and/or human MAGUIN-2, or said fragments, or derivatives, or variants thereof.

20. A method of testing a compound, preferably of screening a plurality of compounds, to determine the degree of binding of said compound or compounds to human MAGUIN-1 and/or human MAGUIN-2, or to fragments, or derivatives, or variants thereof, said method comprising the steps of:

- (i) adding a liquid suspension of said human MAGUIN-1 and/or human MAGUIN-2, or fragments, or derivatives, or variants thereof, to a plurality of containers;
- (ii) adding a detectable compound, preferably a plurality of detectable compounds, in particular fluorescently detectable compounds, to be screened for said binding to said plurality of containers;
- (iii) incubating the liquid suspension of said human MAGUIN-1 and/or human MAGUIN-2, or said fragments, or derivatives, or variants thereof, and said compound, preferably said plurality of compounds;
- (iv) measuring amounts of detectable compound or fluorescence associated with said human MAGUIN-1 and/or human MAGUIN-2, or with said fragments, or derivatives, or variants thereof; and
- (v) determining the degree of binding by one or more of said compounds to said human MAGUIN-1 and/or human MAGUIN-2, or said fragments, or derivatives, or variants thereof.

21. A method for producing a medicament comprising the steps of (i) identifying a modulator of neurodegenerative diseases, in particular Alzheimer's disease, by a method according to claim 18 and (ii) admixing the modulator with a pharmaceutical carrier.

22. A method for producing a medicament comprising the steps of (i) identifying a compound as an inhibitor of binding between a ligand and a human MAGUIN-1 and/or human MAGUIN-2 gene product by a method according to claim 19 and (ii) admixing the compound with a pharmaceutical carrier.

23. A method for producing a medicament comprising the steps of (i) identifying a compound as a binder to a human MAGUIN-1 and/or human MAGUIN-2 gene product by a method according to claim 20 and (ii) admixing the compound with a pharmaceutical carrier.

24. A medicament obtainable by any of the methods according to claim 21 to 23.

25. A medicament obtained by any of the methods according to claim 21 to 23.

26. A protein molecule shown in SEQ ID NO.1 or SEQ ID NO.2, or a fragment, or derivative, or variant thereof, for use as a diagnostic target for detecting a neurodegenerative disease, preferably Alzheimer's disease.

27. A protein molecule shown in SEQ ID NO. 1 or SEQ ID NO.2, or a fragment, or derivative, or variant thereof, for use as a screening target for reagents or compounds preventing, or treating, or ameliorating a neurodegenerative disease, preferably Alzheimer's disease.

28. An antibody specifically immunoreactive with an immunogen, wherein said immunogen is a protein molecule shown in SEQ ID NO. 1, or a fragment, or derivative, or variant thereof.

29. An antibody specifically immunoreactive with an immunogen, wherein said immunogen is a protein molecule shown in SEQ ID NO. 2, or a fragment, or derivative, or variant thereof.

30. Use of an antibody of claim 28 or 29, for detecting the pathological state of a cell in a sample from a subject, comprising immunocytochemical staining of said cell with said antibody, wherein an altered degree of staining, or an altered staining pattern in said cell compared to a cell representing a known health status indicates a pathological state of said cell.

-1/26-

**Fig. 1: Identification of Genes Involved
in Alzheimer's Disease Pathology**




Fig. 2: Identification of differentially expressed genes in a fluorescence differential display screen




Fig. 3: Verification of differential expression of human MAGUIN-1 by quantitative RT-PCR




Fig. 4: Verification of differential expression of Maguin-1 by quantitative RT-PCR






Fig. 5: Verification of differential expression of human MAGUIN-2 by quantitative RT-PCR



-6/26-

Fig. 6: Verification of differential expression of Maguin-2 by quantitative RT-PCR



-7/26-

**Fig. 7: SEQ ID NO. 1:
amino acid sequence of
human MAGUIN-1 protein**

Length: 1034 aa

1 MALIMEPVSK WSPSQVVDWM KGLDDCLQQY IKNFEREKIS GDQLLRITHQ
51 BLEDLGVSRI GHQELILEAV DLLCALNYGL ETENLKTLSH KLNASAKNLQ
101 NFITGRRRSG HYDGRTSRKL PNDFLTSVVD LIGAAKSLLA WLDRSPFAAV
151 TDYSVTRNNV IQLCLELTTI VQQDCTVYET ENKILHVCKT LSGVCDHIIS
201 LSSDPLVSQS AHLEVIQLAN IKPSEGLGMY IKSTYDGLHV ITGTTENSPA
251 DRCKKIHAGD EVIQVNHQTV VGWQLKNLVN ALREDPSGVI LTLKKRPQSM
301 LTSAPALLKN MRWKPLALQP LIPRSPTSSV ATPSSTISTP TKRDSSALQD
351 LYIPPPPAEP YIPRDEKGNL PCEDLRGHMV GKPVHKGSES PNSFLDQEYR
401 KRFNIVEEDT VLYCYEYEKG RSSSQGRRES TPTYGKLRPI SMPVEYNWVG
451 DYEDPNKMKR DSRRENSLLR YMSNEKIAQE EYMFQRNSKK DTGKKSKKKK
501 DKSNSPTHYS LLPSLQMDAL RQDIMGTPVP ETTLYHTFQQ SSLQHKSKKK
551 NKGPIAGKSK RRISCKDLGR GDCEGWLWKK KDAKSYFSQK WKKYWFVLKD
601 ASLYWYINEE DEKAEGFISL PEFKIDRASE CRKKYAFKAC HPKIKSFYFA
651 AEHLDDMNRW LNRINMLTAG YAERERIKQE QDYWSESDKE EADTPSTPKQ
701 DSPPPPYDTY PRPPSMSCAS PYVEAKHSRL SSTETSQSQS SHEEFRQEV
751 GSSAVSPIRK TASQRRSWQD LIETPLTSSG LHYLQTLPLE DSVFSDSAI
801 SPEHRRQSTL PTQKCHLQDH YGPYPLAESE RMQVLNNGG KPRSFTLPRD
851 SGFNHCCLNA PVSACDPQDD VQPPEVEEEE EEEEEEGEAA GENIGEKSSES
901 REEKLGDSLQ DLYRALEQAS LSPLGEHRIS TKMEYKLSFI KRCNDPVMNE
951 KLHRLRLKS TLKAREGEVA IIDKVLDNPD LTSKEFQQWK QMYLDLFLDI
1001 CQNTTSNDPL SISSEVDVIT SSLAHTHSYI ETHV*

-8/26-

**Fig. 8: Alignment of SEQ ID NO. 1,
human MAGUIN-1, with
rat MAGUIN-1**

Length: 1034 aa

```
1 MALIMEPVSKWSPSQVVDWMKGLDDCLOQYIKNFEREKISGDQLLRITHQ 50
  ||||||||||||||||||||||||||||||||||||||||||||||||
1 MALIMEPVSKWSPSQVVDWMKGLDDCLOQYIKNFEREKISGDQLLRITHQ 50

51 ELEDLGVSRIHQELILEAVDLLCALNYGLETENLKTLSHKLNASAKNLQ 100
  ||||||||||||||||||||||||||||||||||||||||||||||||
51 ELEDLGVSRIHQELILEAVDLLCALNYGLETENLKTLSHKLNASAKNLQ 100

101 NFITGRRRSRGHYDGRTSRKLPNDFLT SVVDLIGA AKSLLAWLDRSPFAAV 150
  ||||||||||||||||||||||||||||||||||||||||||||||||
101 NFITGRRRSRGHYDGRTSRKLPNDFLT SVVDLIGA AKSLLAWLDRSPFAAV 150

151 TDYSVTRNNVIQLCLELTTIVQQDCTVYETENKILHVCKTSLSGVCDHIIS 200
  ||||||||||||||||||||||||||||||||||||||||||||||||
151 TDYSVTRNNVIQLCLELTTIVQQDCTVYETENKILHVCKTSLSGVCDHIIS 200

201 LSSDPLVSQSAHLEVIQLANIKPSEGLGMYIKSTYDGLHVITGTTENSPA 250
  ||||||||||||||||||||||||||||||||||||||||||||||||
201 LSSDPLVSQSAHLEVIQLANIKPSEGLGMYIKSTYDGLHVITGTTENSPA 250

251 DRCKKIHAGDEVIQVNHQTVVGWQLKNLVNALREDPSGVILTLKKRPQSM 300
  ||||||||||||||||||||||||||||||||||||||||||||||||
251 DRCKKIHAGDEVIQVNHQTVVGWQLKNLVNALREDPSGVILTLKKRPQSM 300

301 LTSAPALLKNMRWKPLALQPLIPRSPTSSVATPSSTISTPTKRDSSALQD 350
  ||||||||||||||||||||||||||||||||||||||||||||||||
301 LTSAPALLKNMRWKPLALQPLIPRSPTSSVATPSSTISTPTKRDSSALQD 350

351 LYIPPPPAEPYIPRDEKGNLPCEDLRGHMVGKPVHKGSESPNSFLDQEYR 400
  ||||||||||||||||||||||||||||||||||||||||||||||||
351 LYIPPPPAEPYIPRDEKGNLPCEDLRGHMVGKPVHKGSESPNSFLDQEYR 400

401 KRFNIVEEDTVLYCYEYEKGRSSSQGRRESTPTYGKL RPISMPVEYNWVG 450
  ||||||||||||||||||||||||||||||||||||||||||||||||
401 KRFNIVEEDTVLYCYEYEKGRSSSQGRRESTPTYGKL RPISMPVEYNWVG 450

451 DYEDPNKMKRDSRRENSLLRYMSNEKIAQEEYMFQRNSK KDTGKKS KKKG 500
  ||||||||||||||||||||||||||||||||||||||||||||||||
451 DYEDPNKMKRDSRRENSLLRYMSNEKIAQEEYMFQRNSK KDTGKKS KKKG 500
```

-9/26-

501 DKSNSPTHYSLPSLQMDALRQDIMGTPVPETTLYHTFQQSSLQHKSKKK 550
|||:||||||||||||||||||||||||||||||||||||||||||
501 DKSTSPTHYSLPSLQMDALRQDIMGTPVPETTLYHTFQQSSLQHKSKKK 550
551 NKGPIAGKSKRRISCKDLGRGDCEGWLWKKKDAKSYFSQKWKKYWFVLKD 600
|||:||||||||||||||||||||||||||||||||||||||||||
551 NKGAIAGKSKRRISCKDLGRGDCEGWLWKKKDAKSYFSQKWKKYWFVLKD 600
601 ASLYWYINEEDEKAEGFISLPEFKIDRASECRKKYAFKACHPKIKSFYFA 650
||||||||||||||||||||||||||||||||||||||||||
601 ASLYWYINEEDEKAEGFISLPEFKIDRASECRKKYAFKACHPKIKSFYFA 650
651 AEHLDDMNRWLNLRINMLTAGYAERERIKQEODYWSESDKEEADTPSTPKQ 700
||||||||||||||||||||||||||||||||||||||||||
651 AEHLDDMNRWLNLRINMLTAGYAERERIKQEODYWSESDKEEADTPSTPKQ 700
701 DSPPPPYDTPRPPSMSCASPYVEAKHSRLSSTETSQSQSSHEEFRQEV 750
||||||||||||||||||||||||||||||||||||||||||
701 DSPPPPYDTPRPPSMSCASPYVEAKHSRLSSTETSQSQSSHEEFRQEV 750
751 GSSAVSPIRK TASQRRSWQDLIETPLTSSGLHYLQTLPLEDSVFSDSA 800
||||||||||||||||||||||||||||||||||||||||||
751 GSSAVSPIRK TASQRRSWQDLIETPLTSSGLHYLQTLPLEDSVFSDSA 800
801 SPEHRRQSTLPTQKCHLQDHYGPYPLAESERMQVLNGNGGKPRSFTLPRD 850
||||||||||||||||||||||||||||||||||||||||||
801 SPEHRRQSTLPTQKCHLQDHYGPYPLAESERMQVLNGNGGKPRSFTLPRD 850
851 SGFNHCCLNAPVSACDPQDDVQPPEVEEEEEEEEEEGEAAGENIGEKSSES 900
||||||||||||||||||:|||||||||||||||:|:|:|
851 SGFNHCCLNAPVSACDPQDDIQPPEVEEEEEEEEE..EAAGENIGEKNEN 898
901 REEKLGD SLQDLYRALEQASLSPLGEHRISTKMEYKLSFIKRCNDPVMNE 950
||||||||||||||||||:|||||||||||||||:|:|:|
899 REEKLGD SLQDLYRALEEASLSPLGEHRISTKIEYKLSFIKRCNDPVMNE 948
951 KLHRLRILKSTLKAREGEVAIIDKVLDNPDLT SKEFQQWKQMYLDLFLDI 1000
||||||||||||||||||||||||||||||||||||||||||
949 KLHRLRILKSTLKAREGEVAIIDKVLDNPDLT SKEFQQWKQMYLDLFLDI 998
1001 CQNTTSNDPLSISSEVDVITSSLAHTHSYIETHV 1034
||||||||||||||||||:|||||||||||
999 CQNTTSNDPLSISSEVDVITSSLAHTHSYIETHV 1032

-10/26-

Fig. 9: SEQ ID NO. 2: amino acid sequence of human MAGUIN-2 protein

Length: 898 aa

1 MALIMEPVSK WSPSQVVDWM KGLDDCLQQY IKNFEREKIS GDQLLRITHQ
51 ELEDLGVSRI GHQELILEAV DLLCALNYGL ETENLKTLSH KLNASAKNLQ
101 NFITGRRRSG HYDGRTSRKL PNDFLTSVVD LIGAAKSLA WLDRSPFAAV
151 TDYSVTRNNV IQLCLELTTI VQQDCTVYET ENKILHVCKT LSGVCDHIIS
201 LSSDPLVSQS AHLEVIQLAN IKPSEGLGMY IKSTYDGLHV ITGTTENSPA
251 DRCKKIHAGD EVIQVNHQTV VGWQLKNLVN ALREDPSGVI LTLKKRPQSM
301 LTSAPALLKN MRWKPLALQP LIPRSPTSSV ATPSSTISTP TKRDSSALQD
351 LYIPPPPAEP YIPRDEKGNL PCEDLRGHMV GKPVHKGSES PNSFLDQEYR
401 KRFNIVEEDT VLYCYEYEKG RSSSQGRRES TPTYGKLRPI SMPVEYNWVG
451 DYEDPNKMKR DSRRENSLLR YMSNEKIAQE EYMFQRNSKK DTGKKSKKKG
501 DKSNSPTHYS LLPSLQMDAL RQDIMGTPVP ETTLYHTFQQ SSLQHKSKKK
551 NKGPIAGKSK RRISCKDLGR GDCEGWLWKK KDAKSYFSQK WKKYWFLKD
601 ASLYWYINEE DEKAEGFISL PEFKIDRASE CRKKYAFKAC HPKIKSFYFA
651 AEHLDDMNRW LNRINMLTAG YAERERIKQE QDYWSESDKE EADTPSTPKQ
701 DSPPPPYDTY PRPPSMSCAS PYVEAKHSRL SSTETSQSQS SHEEFRQEV
751 GSSAVSPIRK TASQRRSWQD LIETPLTSSG LHYLQTLPLE DSVFSDSAI
801 SPEHRRQSTL PTQKCHLQDH YGPYPLAESE RMQVLNGNGG KPRSFTLPRD
851 SGFNHCCLNA PVSACDPQDD VQPPEVEEEE EEEEEGEAA GENIGEKS*

-11/26-

**Figure 10: Alignment of SEQ ID NO. 2,
human MAGUIN-2, with
rat MAGUIN-2**

Length: 898 aa

```
1 MALIMEPVSKWSPSQVVDWMKGLDDCLQQYIKNFEREKISGDQLLRITHQ 50
  |||
1 MALIMEPVSKWSPSQVVDWMKGLDDCLQQYIKNFEREKISGDQLLRITHQ 50

51 ELEDLGVSRIHQELILEAVDLLCALNYGLETENLKTLSHKLNASAKNLQ 100
  |||
51 ELEDLGVSRIHQELILEAVDLLCALNYGLETENLKTLSHKLNASAKNLQ 100

101 NFITGRRRSRGHYDGRTSRKLPNDFLT SVVDLIGAAKSL LAWLDRSPFAAV 150
  |||
101 NFITGRRRSRGHYDGRTSRKLPNDFLT SVVDLIGAAKSL LAWLDRSPFAAV 150

151 TDYSVTRNNVIQLCLELTTIVQQDCTVYETENKILHVCKTSLSGVCDHIIS 200
  |||
151 TDYSVTRNNVIQLCLELTTIVQQDCTVYETENKILHVCKTSLSGVCDHIIS 200

201 LSSDPLVSQSAHLEVIQLANIKPSEGLGMYIKSTYDGLHVITGTTENSPA 250
  |||
201 LSSDPLVSQSAHLEVIQLANIKPSEGLGMYIKSTYDGLHVITGTTENSPA 250

251 DRCKKIHAGDEVIQVNHQTVVGWQLKNLVNALREDPSGVILTLKKRPQSM 300
  |||
251 DRCKKIHAGDEVIQVNHQTVVGWQLKNLVNALREDPSGVILTLKKRPQSM 300

301 LTSAPALLKNMRWKPLALQPLIPRSPTSSVATPSSTISTPTKRDSSALQD 350
  |||
301 LTSAPALLKNMRWKPLALQPLIPRSPTSSVATPSSTISTPTKRDSSALQD 350

351 LYIPPPPAEPYIPRDEKGNLPCEDLRGHMVGKPVHKGSESPNSFLDQEYR 400
  |||
351 LYIPPPPAEPYIPRDEKGNLPCEDLRGHMVGKPVHKGSESPNSFLDQEYR 400

401 KRFNIVEEDTVLYCYEYEKGRSSSQGRRESTPTYGKLRPISMPVEYNWVG 450
  |||
401 KRFNIVEEDTVLYCYEYEKGRSSSQGRRESTPTYGKLRPISMPVEYNWVG 450

451 DYEDPNKMKRDSRRENSLLRYMSNEKIAQE EYMFQRNSKKDTGKKS KKKG 500
  |||
451 DYEDPNKMKRDSRRENSLLRYMSNEKIAQE EYMFQRNSKKDTGKKS KKKG 500
```

-12/26-

501 DKSNSPTHYSLPSLQMDALRQDIMGTPVPETTLYHTFQQSSLQHKS KKK 550
|||:|||||||||||||||||||||||||||||||||||||
501 DKSTSPTHYSLPSLQMDALRQDIMGTPVPETTLYHTFQQSSLQHKS KKK 550
551 NKGPIAGKSKRRISCKDLGRGDCEGWLWKKKDAKSYFSQKWKKYWFVLKD 600
|||:|||||||||||||||||||||||||||||||||||||
551 NKGAIAGKSKRRISCKDLGRGDCEGWLWKKKDAKSYFSQKWKKYWFVLKD 600
601 ASLYWYINEEDEKAEGFISLPEFKIDRASECRKKYAFKACHPKIKS FYFA 650
|||||||||||||||||||||||||||||||||||||
601 ASLYWYINEEDEKAEGFISLPEFKIDRASECRKKYAFKACHPKIKS FYFA 650
651 AEHLDDMNRLNRLNMLTAGYAERERIKQEQDYWSESDKEEADTPSTPKQ 700
|||||||||||||||||||||||||||||||||||||
651 AEHLDDMNRLNRLNMLTAGYAERERIKQEQDYWSESDKEEADTPSTPKQ 700
701 DSPPPPPYDTYPRPPSMSCASPYVEAKHSRLSSTETSQSQSSHEEF RQEV 750
|||||||||||||||||||||||||||||||||||||
701 DSPPPPPYDTYPRPPSMSCASPYVEAKHSRLSSTETSQSQSSHEEF RQEV 750
751 GSSAVSPIRKTASQRRSWQDLIETPLTSSGLHYLQTLPLEDSVFSDSAAI 800
|||||||||||||||||||||||||||||||||||||
751 GSSAVSPIRKTASQRRSWQDLIETPLTSSGLHYLQTLPLEDSVFSDSAAI 800
801 SPEHRRQSTLPTQKCHLQDHYGPYPLAESERMQVLNNGGKPRSFTLPRD 850
|||||||||||||||||||||||||||||||||||||
801 SPEHRRQSTLPTQKCHLQDHYGPYPLAESERMQVLNNGGKPRSFTLPRD 850
851 SGFNHCCLNAPVSACDPQDDVQPPEVEEEEEEEEEEGEAAGENIGEKS 898
|||||||||||||||||||:||||||||| |||||
851 SGFNHCCLNAPVSACDPQDDIQPPEVEEEEEEEEE..EAAGENIGEKS 896

-13/26-

Fig. 11: SEQ ID NO. 3: nucleotide sequence of human MAGUIN-1 coding sequence**Length: 3105 bp**

```

1  ATGGCTCTGA  TAATGGAACC  GGTGAGCAAA  TGGTCTCCGA  GTCAAGTAGT  GGAAGTGGATG
61  AAAGGTCTTG  ATGACTGTTT  GCAGCAGTAT  ATTAAGAACT  TTGAGAGGGA  GAAGATCAGT
121  GGGGACCAGC  TGCTGCGCAT  TACACATCAG  GAGCTAGAAG  ATCTGGGGGT  CAGCCGCATT
181  GGCCATCAGG  AACTGATCTT  GGAAGCAGTT  GACCTTCTGT  GTGCATTGAA  TTATGGCTTG
241  GAAACAGAAA  ATCTAAAAAC  CCTTCTCACC  AAGTTGAATG  CATCTGCCAA  AAATCTGCAG
301  AATTTTATAA  CAGGAAGGAG  AAGGAGTGCG  CATTATGATG  GGAGGACCAG  CCGAAAATTG
361  CCAAACGACT  TTCTGACCTC  AGTTGTGGAT  CTGATTGGAG  CAGCCAAGAG  TCTGCTTGCC
421  TGGTTGGACA  GGTCACCATT  TGCTGCTGTG  ACAGACTATT  CAGTTACAAG  AAATAATGTC
481  ATACAACTCT  GCCTGGAGTT  AACAACAATT  GTGCAACAGG  ATTGTACTGT  ATATGAAACA
541  GAGAATAAAA  TTCTTCACGT  GTGTAAAAC  CTTTCTGGAG  TCTGTGACCA  CATCATATCC
601  CTGTCGTCAG  ATCCTCTGGT  TTCACAGTCT  GCTCACCTGG  AAGTGATTCA  ACTGGCAAAC
661  ATTAACCAA  GCGAAGGGCT  GGGTATGTAT  ATTAATCTA  CATATGATGG  CCTCCATGTA
721  ATTAAGTGG  CCACAGAAAA  TTCACCTGCA  GATCGGTGCA  AGAAAATCCA  TGCTGGCGAT
781  GAAGTGATTC  AAGTTAATCA  TCAGACTGTG  TGGTGTATC  TTAACCTTGA  AAAAGCGACC  TCAGAGCATG
841  GCACTACGAG  AGGACCCGAG  TGGTGTATC  TTAACCTTGA  AAAAGCGACC  TCAGAGCATG
901  CTTACCTCAG  CACCAGCTTT  ACTGAAAAAT  ATGAGATGGA  AGCCCTTTCG  TCTGCAGCCT
961  CTTATACCTA  GAAGTCCAC  AAGCAGCGTT  GCCACGCCTT  CCAGCACCAT  CAGTACACCC
1021  ACCAAAAGAG  ACAGTTCTGC  CCTCCAGGAT  CTCTACATTC  CCCCTCCTCC  TGCAGAACCA
1081  TATATTCCCA  GGGATGAAAA  AGGAAACCTT  CCTTGTGAAG  ACCTCAGAGG  ACATATGGTG
1141  GGCAAGCCAG  TGCATAAGGG  ATCTGAATCA  CCAAATTCAT  TTCTGGATCA  GGAATATCGA
1201  AAGAGATTTA  ATATTGTGCA  AGAAGATACT  GTCTTATATT  GCTATGAATA  TGAAAAAGGA
1261  AGATCAAGTA  GTCAAGGAAG  ACGAGAAAGC  ACCCAACTT  ATGGCAAGCT  ACGACCTATA
1321  TCTATGCCAG  TGGAATATA  TTGGTGGGG  GACTATGAAG  ATCCAAATA  GATGAAGAGA
1381  GATAGTAGAA  GAGAAAACTC  TCTACTTCGG  TATATGAGCA  ATGAAAAGAT  TGCTCAAGAA
1441  GAATACATGT  TTCAGAGAAA  CAGCAAAAAG  GACACAGGGA  AGAAGTCAA  AAAGAAGGGT
1501  GATAAGAGTA  ATAGCCCAAC  TCACTATTCA  TTGCTACCTA  GTTTACAAAT  GGATGCACTG
1561  AGACAAGACA  TCATGGGCAC  TCCTGTGCCA  GAGACCACAC  TATACCATAC  ATTTCAGCAG
1621  TCCTCACTGC  AGCACAATC  AAAGAAGAAA  AACAAAGGTC  CTATAGCAGG  CAAGAGCAAA
1681  AGACGAATTT  CTTGCAAAGA  TCTTGGCCGT  GGTGACTGTG  AGGGCTGGCT  TTGGAAGGAT
1741  AAAGATGCGA  AGAGTTACTT  TTCACAGAAA  TGGAAGGAT  ATTGGTTTGT  CTTAAAGGAT
1801  GCATCCCTTT  ATTGGTATAT  TAATGAGGAG  GATGAAAAAG  CAGAAGGATT  CATTAGCCTG
1861  CCTGAATTTA  AAATTGATAG  AGCCAGTGAA  TGCCGCAAAA  AATATGCATT  CAAAGCCTGT
1921  CATCCTAAAA  TCAAAAGCTT  TTATTTTGCT  GCTGAACATC  TTGATGATAT  GAACAGGTGG
1981  CTTAACAGAA  TTAATATGCT  GACTGCAGGA  TATGCAGAAA  GAGAGAGGAT  TAAGCAGGAA
2041  CAAGATTACT  GGAGTGAGAG  TGACAAGGAA  GAAGCAGATA  CTCCATCAAC  ACCAAAACAA
2101  GATAGCCCTC  CACCCCATTA  TGATACATAC  CCACGACCTC  CCTCGATGAG  TTGCGCCAGT
2161  CCTTATGTGG  AAGCAAAACA  TAGCCGACTT  TCCTCCACGG  AGACTTCTCA  GTCTCAGTCT
2221  TCTCATGAGG  AGTTTCGCCA  GGAAGTAACT  GGGAGCAGTG  CAGTGTCTCC  CATTGCAAG
2281  ACAGCCAGTC  AGCGCCGCTC  CTGGCAGGAT  TTAATTGAGA  CGCCACTGAC  AAGTTCAGGC
2341  TTACACTATC  TTCAGACTCT  GCCCCTGGAG  GATTCTGTCT  TCTCTGACTC  CGCGGCCATC
2401  TCCCCAGAGC  ACAGGCGGCA  GTCTACCTG  CCAACTCAGA  AATGCCACCT  GCAGGATCAC
2461  TATGGGCCAT  ACCCCTTAGC  TGAGAGTGAG  AGGATGCAAG  TGCTAAATGG  AAATGGGGGC
2521  AAGCCTCGAA  GTTTTACTCT  GCCTCGAGAT  AGCGGGTTCA  ACCATTGCTG  TCTGAATGCT
2581  CCAGTTAGTG  CCTGTGACCC  ACAGGATGAC  GTGCAACCCC  CAGAGGTGGA  GGAAGAGGAG
2641  GAGGAGGAGG  AGGAGGAAGG  GGAGGCAGCA  GGGGAAAACA  TAGGAGAAAA  AAGTGAAAGC
2701  AGAGAAGAAA  AGTTAGGAGA  CTCATTGCAA  GATTTATACA  GGGCACTGGA  GCAGGCCAGT
2761  CTGTCAACCAC  TAGGAGAACA  TCGTATTTCA  ACCAAGATGG  AATACAAGCT  ATCATTTATA
2821  AAAAGATGTA  ATGATCCTGT  AATGAATGAA  AAACACAC  GGCTGAGAA  TCTCAAAAGC
2881  ACTTTAAAGG  CCAGAGAAGG  GGAAGTAGCC  ATTATCGATA  AAGTCCTAGA  CAATCCAGAC
2941  TTGACATCTA  AAGAATTCCA  ACAATGGAAG  CAGATGTACC  TCGACCTTTT  CTTGGATATC
3001  TGTCAAAATA  CCACCTCAA  TGACCCACTG  AGTATTTCTT  CTGAAGTAGA  TGTAATCACT
3061  TCCTCTCTAG  CACACACTCA  TTCATACATT  GAAACGCATG  TCTAA

```

-14/26-

Fig. 12: SEQ ID NO. 4: nucleotide sequence of human MAGUIN-2 coding sequence

Length: 2697 bp

```

1  ATGGCTCTGA TAATGGAACC GGTGAGCAAA TGGTCTCCGA GTCAAGTAGT
51  GGACTGGATG AAAGGTCTTG ATGACTGTTT GCAGCAGTAT ATTAAGAACT
101 TTGAGAGGGA GAAGATCAGT GGGGACCAGC TGCTGCGCAT TACACATCAG
151 GAGCTAGAAG ATCTGGGGGT CAGCCGCATT GGCCATCAGG AACTGATCTT
201 GGAAGCAGTT GACCTTCTGT GTGCATTGAA TTATGGCTTG GAAACAGAAA
251 ATCTAAAAAC CCTTTCTCAC AAGTTGAATG CATCTGCCAA AAATCTGCAG
301 AATTTTATAA CAGGAAGGAG AAGGAGTGGC CATTATGATG GGAGGACCAG
351 CCGAAAATTG CCAAACGACT TTCTGACCTC AGTTGTGGAT CTGATTGGAG
401 CAGCCAAGAG TCTGCTTGCC TGGTTGGACA GGTCACCATT TGCTGCTGTG
451 ACAGACTATT CAGTTACAAG AAATAATGTC ATACAACTCT GCCTGGAGTT
501 AACACAATT GTGCAACAGG ATTGTACTGT ATATGAAACA GAGAATAAAA
551 TTCTTCACGT GTGTAAACT CTTTCTGGAG TCTGTGACCA CATCATATCC
601 CTGTCGTCAG ATCCTCTGGT TTCACAGTCT GCTCACCTGG AAGTGATTCA
651 GCTGGCAAAC ATTAAACCAA GCGAAGGGCT GGGTATGTAT ATTAAATCTA
701 CATATGATGG CCTCCATGTA ATTACTGGAA CCACAGAAAA TTCACCTGCA
751 GATCGGTGCA AGAAAATCCA TGCTGGCGAT GAAGTGATTG AAGTTAATCA
801 TCAGACTGTG GTGGGGTGGC AGTTGAAAAA TTTGGTGAAT GCACTACGAG
851 AGGACCCGAG TGGTGTATC TTAACCTTGA AAAAGCGACC TCAGAGCATG
901 CTTACCTCAG CACCAGCTTT ACTGAAAAAT ATGAGATGGA AGCCCCCTGC
951 TCTGCAGCCT CTTATACCTA GAAGTCCAC AAGCAGCGTT GCCACGCCTT
1001 CCAGCACCAT CAGTACACCC ACCAAAAGAG ACAGTTCTGC CCTCCAGGAT
1051 CTCTACATTC CCCCTCCTCC TGCAGAACCA TATATTCCCA GGGATGAAAA
1101 AGGAAACCTT CCTTGTAAG ACCTCAGAGG ACATATGGTG GGCAAGCCAG
1151 TGCATAAGGG ATCTGAATCA CCAAATTCAT TTCTGGATCA GGAATATCGA
1201 AAGAGATTTA ATATTGTCTGA AGAAGATACT GTCTTATATT GCTATGAATA
1251 TGAAAAAGGA AGATCAAGTA GTCAAGGAAG ACGAGAAAGC ACCCCAACCTT
1301 ATGGCAAGCT ACGACCTATA TCTATGCCAG TGGAATATAA TTGGGTGGGG
1351 GACTATGAAG ATCCAAATAA GATGAAGAGA GATAGTAGAA GAGAAAACTC
1401 TCTACTCGG TATATGAGCA ATGAAAAGAT TGCTCAAGAA GAATACATGT
1451 TTCAGAGAAA CAGCAAAAAG GACACAGGGA AGAAGTCAA AAAGAAGGGT
1501 GATAAGAGTA ATAGCCCAAC TCACTATTCA TTGCTACCTA GTTTACAAAT
1551 GGATGCACTG AGACAAGACA TCATGGGCAC TCCTGTGCCA GAGACCACAC
1601 TATACCATAC ATTTTCAGCAG TCCTCACTGC AGCACAAATC AAAGAAGAAA
1651 AACAAAGGTC CTATAGCAGG CAAGAGCAAA AGACGAATTT CTTGCAAAGA
1701 TCTTGCCCGT GGTGACTGTG AGGGCTGGCT TTGGA AAAAG AAAGATGCGA
1751 AGAGTTACTT TTCACAGAAA TGGAAAAAAT ATTGGTTTGT CCTAAAGGAT
1801 GCATCCCTTT ATTGGTATAT TAATGAGGAG GATGAAAAAG CAGAAGGATT
1851 CATTAGCCTG CCTGAATTTA AAATTGATAG AGCCAGTGAA TGCCGCAAAA
1901 AATATGCATT CAAAGCCTGT CATCCTAAAA TCAAAAGCTT TTATTTTGCT
1951 GCTGAACATC TTGATGATAT GAACAGGTGG CTTAACAGAA TTAATATGCT
2001 GACTGCAGGA TATGCAGAAA GAGAGAGGAT TAAGCAGGAA CAAGATTACT
2051 GGAGTGAGAG TGACAAGGAA GAAGCAGATA CTCCATCAAC ACCAAAACAA
2101 GATAGCCCTC CACCCCATTA TGATACATAC CCACGACCTC CCTCGATGAG
2151 TTGCGCCAGT CCTTATGTGG AAGCAAAACA TAGCCGACTT TCCTCCACGG
2201 AGACTTCTCA GTCTCAGTCT TCTCATGAGG AGTTTCGCCA GGAAGTAACT
2251 GGGAGCAGTG CAGTGTCTCC CATTCGCAAG ACAGCCAGTC AGCGCCGCTC
2301 CTGGCAGGAT TTAATTGAGA CGCCACTGAC AAGTTCAGGC TTACACTATC
2351 TTCAGACTCT GCCCCTGGAG GATTCTGTCT TCTCTGACTC CGCGGCCATC
2401 TCCCAGAGC ACAGGCGGCA GTCTACCCTG CCAACTCAGA AATGCCACCT
2451 GCAGGATCAC TATGGGCCAT ACCCCTTAGC TGAGAGTGAG AGGATGCAAG
2501 TGCTAAATGG AAATGGGGGC AAGCCTCGAA GTTTTACTCT GCCTCGAGAT
2551 AGCGGGTTCA ACCATTGCTG TCTGAaTGCT CCAGTTAGTG CCTGTGACCC
2601 ACAGGATGAC GTGCAACCCC CAGAGGTGGA GGAAGAGGAG GAGGAGGAGG
2651 AGGAGGAAGG GGAGGCAGCA GGGGAAAACA TAGGAGAAAA AAGCTAA

```

-15/26-**Fig. 13: SEQ ID NO. 5: nucleotide sequence of human MAGUIN-1 cDNA****Length: 5749 bp**

```
1  CGGGCAGCTA  GTCGTGCTCG  GGGCTTCACT  CCCGCGCGTG  AGGCGAGCGG  GCAAGTTGGC
61  TGAGGGCGTG  CGGCAGAGGC  TGCTTCCCTC  GGCGACGCGA  CCCCTCAGCA  ACTCAAGCTA
121  TGAACGAAAG  CTCCCTAGGG  ACGGAGACCG  GAGCGGAGCG  GCGGAGGCAG  CAGCAGCAGC
181  AGCAGCAGCA  GCAGCAGCAG  CAGCCGCCGC  CGCCGCCGCC  TTAGCGGGAA  CTGAGCAGAC
241  CCGGCGCGGA  GCCACGACTC  CTGCACGTTT  ACCTCCCTGT  CGCCGTTCCT  GCCGGCGGTT
301  GGCTAAAAGA  CGTTACAGCC  GCGAGACCCG  ACACACAAAA  GCCGCTTTC  CCGCGCCGCC
361  CGCCAGGGA  GGCTGCGGCC  AGCAAGGGAC  CCCACCTGAG  AGCAGCTCGG  GCTGCTGAGT
421  TCGTTTTGTG  TCTGAGCTCT  GCGCTCTGCA  CGGAACCGAC  CCCGTACCCA  TGGCTCTGAT
481  AATGGAACCG  GTGAGCAAAT  GGTCTCCGAG  TCAAGTAGTG  GACTGGATGA  AAGGTCTTGA
541  TGACTGTTTG  CAGCAGTATA  TTAAGAACTT  TGAGAGGGAG  AAGATCAGTG  GGGACCAGCT
601  GCTGCGCATT  ACACATCAGG  AGCTAGAAGA  TCTGGGGGTC  AGCCGCATTG  GCCATCAGGA
661  ACTGATCTTG  GAAGCAGTTG  ACCTTCTGTG  TGCATTGAAT  TATGGCTTGG  AAACAGAAAA
721  TCTAAAACC  CTTTCTCACA  AGTTGAATGC  ATCTGCCAAA  AATCTGCAGA  ATTTTATAAC
781  AGGAAGGAGA  AGGAGTGGCC  ATTATGATGG  GAGGACCAGC  CGAAAATTGC  CAAACGACTT
841  TCTGACCTCA  GTTGTGGATC  TGATTGGAGC  AGCCAAGAGT  CTGCTTGCCT  GGTTGGACAG
901  GTCACCATT  GCTGCTGTGA  CAGACTATTC  AGTTACAAGA  AATAATGTCA  TACAACCTG
961  CCTGGAGTTA  ACAACAATTG  TGCAACAGGA  TTGTACTGTA  TATGAAACAG  AGAATAAAAT
1021  TCTTCACGTG  TGTAAAACTC  TTTCTGGAGT  CTGTGACCAC  ATCATATCCC  TGTCTCAGCA
1081  TCCTCTGGTT  TCACAGTCTG  CTCACCTGGA  AGTGATTCAA  CTGGCAAACA  TTAAACCAAG
1141  CGAAGGGCTG  GGTATGTATA  TTAATCTTAC  ATATGATGGC  CTCCATGTAA  TTACTGGAAC
1201  CACAGAAAT  TCACCTGCAG  ATCGGTGCAA  GAAATCCAT  GCTGGCGATG  AAGTGATTCA
1261  AGTTAATCAT  CAGACTGTGG  TGGGGTGGCA  GTTGAAAAAT  TTGGTGAATG  CACTACGAGA
1321  GGACCCGAGT  GGTGTTATCT  TAACTTTGAA  AAAGCGACCT  CAGAGCATGC  TTACCTCAGC
1381  ACCAGCTTTA  CTGAAAAATA  TGAGATGGAA  GCCCCTTGCT  CTGCAGCCTC  TTATACCTAG
1441  AAGTCCCACA  AGCAGCGTTG  CCACGCCTTC  CAGCACCATC  AGTACACCCA  CCAAAAGAGA
1501  CAGTTCTGCC  CTCCAGGATC  TCTACATTCC  CCTCCTCCT  GCAGAACCAT  ATATTCCAG
1561  GGATGAAAAA  GGAAACCTTC  CTTGTGAAGA  CCTCAGAGGA  CATATGGTGG  GCAAGCCAGT
1621  GCATAAGGGA  TCTGAATCAC  CAAATTCATT  TCTGGATCAG  GAATATCGAA  AGAGATTTAA
1681  TATTGTCCGA  GAAGATACTG  TCTTATATTG  CTATGAATAT  GAAAAAGGAA  GATCAAGTAG
1741  TCAAGGAAGA  CGAGAAAAGCA  CCCCACCTTA  TGGCAAGCTA  CGACCTATAT  CTATGCCAGT
1801  GGAATATAAT  TGGGTGGGGG  ACTATGAAGA  TCCAAATAAG  ATGAAGAGAG  ATAGTAGAAG
1861  AGAAAACCTC  CTACTTCGGT  ATATGAGCAA  TGAAAAGATT  GCTCAAGAAG  AATACATGTT
1921  TCAGAGAAAC  AGCAAAAAGG  ACACAGGGAA  GAAGTCAAAA  AAGAAGGGTG  ATAAGAGTAA
1981  TAGCCCAACT  CACTATTTCAT  TGCTACCTAG  TTTACAAATG  GATGCACTGA  GACAAGACAT
2041  CATGGGCACT  CCTGTGCCAG  AGACCACACT  ATACCATACA  TTTCAGCAGT  CCTCACTGCA
2101  GCACAAATCA  AAGAAGAAAA  ACAAAGGTCC  TATAGCAGGC  AAGAGCAAAA  GACGAATTTT
2161  TTGCAAAGAT  CTTGGCCGTG  GTGACTGTGA  GGGCTGGCTT  TGGAAAAAGA  AAGATGCCAA
2221  GAGTTACTTT  TCACAGAAAT  GGAAAAATA  TTGGTTTGTC  CTAAAGGATG  CATCCCTTTA
2281  TTGGTATATT  AATGAGGAGG  ATGAAAAAGC  AGAAGGATTC  ATTAGCCTGC  CTGAATTTAA
2341  AATTGATAGA  GCCAGTGAAT  GCCGCAAAAA  ATATGCATT  AAAGCCTGTC  ATCCTAAAAT
2401  CAAAAGCTTT  TATTTTGCTG  CTGAACATCT  TGATGATATG  AACAGGTGGC  TTAACAGAAT
2461  TAATATGCTG  ACTGCAGGAT  ATGCAGAAAG  AGAGAGGATT  AAGCAGGAAC  AAGATTACTG
2521  GAGTGAGAGT  GACAAGGAAG  AAGCAGATAC  TCCATCAACA  CCAAAACAAG  ATAGCCCTCC
2581  ACCCCCATAT  GATACATACC  CACGACCTCC  CTCGATGAGT  TGCGCCAGTC  CTTATGTGGA
2641  AGCAAAACAT  AGCCGACTTT  CCTCCACGGA  GACTTCTCAG  TCTCAGTCTT  CTCATGAGGA
2701  GTTTCGCCAG  GAAGTAACTG  GGAGCAGTGC  AGTGTCTCCC  ATTCGCAAGA  CAGCCAGTCA
2761  GCGCCGCTCC  TGGCAGGATT  TAATTGAGAC  GCCACTGACA  AGTTCAGGCT  TACACTATCT
2821  TCAGACTCTG  CCCCTGGAGG  ATTCTGTCTT  CTCTGACTCC  GCGGCCATCT  CCCAGAGCA
```


-16/26-

2881 CAGGCGGCAG TCTACCCTGC CAACTCAGAA ATGCCACCTG CAGGATCACT ATGGGCCATA
2941 CCCCTTAGCT GAGAGTGAGA GGATGCAAGT GCTAAATGGA AATGGGGGCA AGCCTCGAAG

3001 TTTTACTCTG CCTCGAGATA GCGGGTTCAA CCATTGCTGT CTGAATGCTC CAGTTAGTGC
3061 CTGTGACCCA CAGGATGACG TGCAACCCCC AGAGGTGGAG GAAGAGGAGG AGGAGGAGGA
3121 GGAGGAAGGG GAGGCAGCAG GGGAAAAACAT AGGAGAAAAA AGTGAAAGCA GAGAAGAAAA
3181 GTTAGGAGAC TCATTGCAAG ATTTATACAG GGCACCTGGAG CAGGCCAGTC TGTCACCACT
3241 AGGAGAACAT CGTATTTCAA CCAAGATGGA ATACAAGCTA TCATTTATAA AAAGATGTAA
3301 TGATCCTGTA ATGAATGAAA AACTACACCG GCTGAGAATT CTCAAAAGCA CTTTAAAGGC
3361 CAGAGAAGGG GAAGTAGCCA TTATCGATAA AGTCCTAGAC AATCCAGACT TGACATCTAA
3421 AGAATTCCAA CAATGGAAGC AGATGTACCT CGACCTTTTC TTGGATATCT GTCAAAATAC
3481 CACCTCAAAT GACCCACTGA GTATTTCTTC TGAAGTAGAT GTAATCACTT CCTCTCTAGC
3541 ACACACTCAT TCATACATTG AAACGCATGT CTAAATGTAT TCTGCCTTCA GACCATCTAG
3601 TACCTGCTGG TACTCTGAAC AAGTATATAA GGTAGTTTTT ATATCAATGT GTGGAACACT
3661 TGACAAGCTA TACTTTAATG TTACCAAACCT ATATGAAACA AACCATATAT GGTCACAATA
3721 CCACTATCTT TAATGAGCAT TTGTATATTT TATATGCAAC AGTGCTCAGC TTATGTTTAC
3781 CATGTGCAAA ATCAACTGTC TTTAATGACT TAAAATTAAC TTTTGCAAAC AATTCTAAAT
3841 ACAGGTGGTC TTCAAGTAGT AAAACCACAA AAGGCAGTTT TCTATCTATG GTCATCTTTT
3901 CTCCCTTTAA GTTAATTTTA TATAACAAG ACTTCAAAG TAAATCACAT TTTTTCAGGT
3961 GCAGACATCC TTGTGGGTGG GAAAGAATTT AAACCTTTTT TATATTTATT AAAATGTTCT
4021 AAGAATTTTC TTAAACATTG CACAAAGTTT AATGCTGTAG TTTTATTTTT GTGAAATGTA
4081 GATGCGCATA CAAGAGCTAA GCAAAATAGA AGAGCATCGA CATAAGAAAA GTTCAGGTAT
4141 CTAATATTCTG TCTTAATAGT CTATTAACCT GTGAAAGCTA AGTTAATGGA AATATTATTC
4201 CAAATCTATG AGAACACTTG GTGTATCAGG GCAAAGCTTT GTAAGATGTT TTTGTAACTA
4261 AGACCAAAAT TGAAGATAGA GCTGCTTTAT TTTCTTGGTT TAAATCTTCC TTTATTTTTG
4321 TAGTGATGAG ATGCTGATTG TGTACAGAAG AATTTGAGAG GGGATTTTTA AAAACTGACT
4381 TAACACACCC AGAAAGGCAG CTAACAGCTA TATATATATA TAAATTTTCA CCCAACTCA
4441 TGTTTTTAAA CTCCAACCTC TAAAAGACAA CAAGGTATAA ACTGAAATGA ATCAACTTTC
4501 CACTTAGTTT CCAATTTTCC CCTAGTCCAC TAATTAACT TAGGTAATTA TACTTCAGGT
4561 AGGGAAGTAC AATATGTTTA GTTTCAGGCT GATGTGTGTT ATAAAAACA ACACTGAAAA
4621 ATAAAAATGT ACTTCCCTTC TAAGGAGCAA GCAGGTGATG GTCATTCAA GAGATGTCAC
4681 ATTGAATTAT GAGAGAAACA ATTTAGAGGT TTTTTCCTG GCTTCATGAA TTGTCTATA
4741 GAGTGGATGA AGTCTAAGGA AAAGTCCTCT TCATATATTT CCATTTATAA GCGTCTTGTT
4801 TTTGAAAGTG ATCACAGCAT GAAAATGACT GTGCTGCTTT TTAGTGCTG GCTGCATAAT
4861 GTACAAAGTCA CAATTTGCTG TTTTTTTCAG GAGGAGAAAG GGAACCTCCT TTACTATTCT
4921 ATATCCTAAA ATCTACTTCT AATCAGCTTT ATACTGTTGC CTGTACAGCT CAGTGAATGT
4981 ACTTTCATCT TTAAGAGTTC AGATATATGC CAGTGAATAT TTTTGCTGTA GAGGAGAAAG
5041 TAAAAACTCC ACAGCGGGGA TCTTTTTCTT TGCTTTTGAA ACCACCATTG AATCACTATC
5101 GTTTTGCAGA CTTTGCACAA CTGTACAGGA GAGTGGCCTT TCTACAGCAC ATTTTCAGTA
5161 ATCCTATATT TAGTCAAAAT GGATGAGAAA TCATGTATTA ATGTTTGTA GGAATTTTGG
5221 GTCCAGTGTA ATATTTTAT CATTTAAAA GAACCTTATT TGTA AAAACA TTTATTTACT
5281 GCATGGATAT TGACGCACAT TAAATTTGTG GGATTTTGTA TATGTAAAAA AAAAAAAAAA
5341 AAAAAAAAAAC AAAAAACCTC TTGTCCTAAA ATGAAGTGTG CTTGTTAACA GGTGTTTAGA
5401 CTTATTGATG TTTACTAGAC CAAATGTGTA TGTTCACTTA AAAATATATG TACCTGATGG
5461 ATGTGTCATG TTTACAGTGG CCAGGTTGTG GCCTGTAAAC AGCAAGCAGT TGACGGGAAG
5521 ACTAGCTCTG TTGCTACTAA GCAGCTTTTA CTTTGTGAAA GTCAGCTCTG TTGTTTTAAA
5581 TGGTAAAAAT TAACTAATG AATTTGACAA GACTCGTGGC TAGCCTAGCA TGAAAGAGAC
5641 CTTTAAACAC TATATAATAT CTGTACATTT TATTGCATTC GTTTCAAATC TAGGAGAGAG
5701 GCAGCACTGT AAACCTGAAGT CAAATAAATT CAGCTCTTAA TGAATCCTT

-17/26-

Fig. 14: SEQ ID NO. 6: nucleotide sequence of human MAGUIN-2 cDNA

Length: 4350 bp

```
1  GTGCTCGGGG CTTCACTCCC GCGCGTGAGG CGAGCGGGCA AGTTGGCTGA
51  GGGCGTGCGG CAGAGGCTGC TTCCCTCGGC GACGCGACCC CTCAGCAACT
101 CAAGCTATGA ACTGAAGCTC CCTAGGGACG GAGACCGGAG CGGAGCGGCG
151 GAGGCAGCAG CAGCAGCAGC AGCAGCAGCA GCAGCAGCAG CCGCCGCCGC
201 CGCCGCCTTA GCGGGAAGTG AGCAGACCCG GCGCGGAGCC ACGACTCCTG
251 CACGTTTACC TCCCTGTCGC CGTTCCTGCC GGCGGTTGGC TAAAAGACGT
301 TACAGCCGCG AGACCCGACA CACAAAAGCC GCTTTCCTCCG CGCCGCCCGC
351 CCAGGGAGGC TGCGGCCAGC AAGGGACCCC ACCTGAGAGC AGCTCGGGCT
401 GCTGAGTTCT TTTTGTGTCT GAGCTCTGCG CTCTGCACGG AACCGACCCC
451 GTACCCATGG CTCTGATAAT GGAACCGGTG AGCAAATGGT CTCCGAGTCA
501 AGTAGTGGAC TGGATGAAAG GTCTTGATGA CTGTTTGCAG CAGTATATTA
551 AGAAGTTTGA GAGGGAGAAG ATCAGTGGGG ACCAGCTGCT GCGCATTACA
601 CATCAGGAGC TAGAAGATCT GGGGGTCAGC CGCATTGGCC ATCAGGAACT
651 GATCTTGGAA GCAGTTGACC TTCTGTGTGC ATTGAATTAT GGCTTGGAAA
701 CAGAAAATCT AAAAACCCCTT TCTCACAAGT TGAATGCATC TGCCAAAAAT
751 CTGCAGAATT TTATAACAGG AAGGAGAAGG AGTGGCCATT ATGATGGGAG
801 GACCAGCCGA AAATTGCCAA ACGACTTTCT GACCTCAGTT GTGGATCTGA
851 TTGGAGCAGC CAAGAGTCTG CTTGCCTGGT TGGACAGGTC ACCATTTGCT
901 GCTGTGACAG ACTATTGAGT TACAAGAAAT AATGTCATAC AACTCTGCCT
951 GGAGTTAACA ACAATTGTGC AACAGGATTG TACTGTATAT GAAACAGAGA
1001 ATAAAATTCT TCACGTGTGT AAAACTCTTT CTGGAGTCTG TGACCACATC
1051 ATATCCCTGT CGTCAGATCC TCTGGTTTCA CAGTCTGCTC ACCTGGAAGT
1101 GATTCAAGCTG GCAAACATTA AACCAAGCGA AGGGCTGGGT ATGTATATTA
1151 AATCTACATA TGATGGCCTC CATGTAATTA CTGGAACCAC AGAAAATTCA
1201 CCTGCAGATC GGTGCAAGAA AATCCATGCT GGCGATGAAG TGATTCAAGT
1251 TAATCATCAG ACTGTGGTGG GGTGGCAGTT GAAAAATTG GTGAATGCAC
1301 TACGAGAGGA CCCGAGTGGT GTTATCTTAA CTTTGAAAAA GCGACCTCAG
1351 AGCATGCTTA CCTCAGCACC AGCTTTACTG AAAAATATGA GATGGAAGCC
1401 CCTTGCTCTG CAGCCTCTTA TACCTAGAAG TCCCACAAGC AGCGTTGCCA
1451 CGCCTTCCAG CACCATCAGT ACACCCACCA AAAGAGACAG TTCTGCCCTC
1501 CAGGATCTCT ACATTCCCCC TCCTCCTGCA GAACCATATA TTCCCAGGGA
1551 TGAAAAAGGA AACCTTCCTT GTGAAGACCT CAGAGGACAT ATGGTGGGCA
1601 AGCCAGTGCA TAAGGGATCT GAATCACCAA ATTCATTTCT GGATCAGGAA
1651 TATCGAAAGA GATTTAATAT TGTGGAAGAA GATACTGTCT TATATTGCTA
1701 TGAATATGAA AAAGGAAGAT CAAGTAGTCA AGGAAGACGA GAAAGCAGCC
1751 CAACTTATGG CAAGCTACGA CCTATATCTA TGCCAGTGGA ATATAATTGG
1801 GTGGGGGACT ATGAAGATCC AAATAAGATG AAGAGAGATA GTAGAAGAGA
1851 AAACCTCTCTA CTTCCGTATA TGAGCAATGA AAAGATTGCT CAAGAAGAAT
1901 ACATGTTTCA GAGAAACAGC AAAAAGGACA CAGGGAAGAA GTCAAAAAAG
1951 AAGGGTGATA AGAGTAATAG CCCAACTCAC TATTCATTGC TACCTAGTTT
2001 ACAAATGGAT GCACTGAGAC AAGACATCAT GGGCACTCCT GTGCCAGAGA
2051 CCACACTATA CCATACATTT CAGCAGTCCT CACTGCAGCA CAAATCAAAG
2101 AAGAAAAACA AAGGTCTTAT AGCAGGCAAG AGCAAAAGAC GAATTTCTTG
2151 CAAAGATCTT GGCCGTGGTG ACTGTGAGGG CTGGCTTTGG AAAAAGAAAG
2201 ATGCGAAGAG TTACTTTTCA CAGAAATGGA AAAAATATTG GTTTGTCCTA
2251 AAGGATGCAT CCCTTTATTG GTATATTAAT GAGGAGGATG AAAAAGCAGA
```

-18/26-

2301 AGGATTCATT AGCCTGCCTG AATTTAAAT TGATAGAGCC AGTGAATGCC
2351 GCAAAAAATA TGCATTCAAA GCCTGTCATC CTAAATCAA AAGCTTTTAT

2401 TTTGCTGCTG AACATCTTGA TGATATGAAC AGGTGGCTTA ACAGAATTAA
2451 TATGCTGACT GCAGGATATG CAGAAAGAGA GAGGATTAAG CAGGAACAAG
2501 ATTACTGGAG TGAGAGTGAC AAGGAAGAAG CAGATACTCC ATCAACACCA
2551 AAACAAGATA GCCCTCCACC CCCATATGAT ACATACCCAC GACCTCCCTC
2601 GATGAGTTGC GCCAGTCCTT ATGTGGAAGC AAAACATAGC CGACTTTCCT
2651 CCACGGAGAC TTCTCAGTCT CAGTCTTCTC ATGAGGAGTT TCGCCAGGAA
2701 GTAACCTGGGA GCAGTGCAGT GTCTCCCATT CGCAAGACAG CCAGTCAGCG
2751 CCGCTCCTGG CAGGATTTAA TTGAGACGCC ACTGACAAGT TCAGGCTTAC
2801 ACTATCTTCA GACTCTGCCC CTGGAGGATT CTGTCTTCTC TGACTCCGCG
2851 GCCATCTCCC CAGAGCACAG GCGGCAGTCT ACCCTGCCAA CTCAGAAATG
2901 CCACCTGCAG GATCACTATG GGCCATACCC CTTAGCTGAG AGTGAGAGGA
2951 TGCAAGTGCT AAATGGAAAT GGGGGCAAGC CTCGAAGTTT TACTCTGCCT
3001 CGAGATAGCG GGTTCACCA TTGCTGTCTG AaTGCTCCAG TTAGTGCCTG
3051 TGACCCACAG GATGACGTGC AACCCCCAGA GGTGGAGGAA GAGGAGGAGG
3101 AGGAGGAGGA GGAAGGGGAG GCAGCAGGGG AAAACATAGG AGAAAAAGC
3151 TAATACACTG CGAGAGTTGG TAGAACCTCT CCATGCCAAA TCGGATCCAC
3201 TTCTGTTGGC ACTCAACCCA TTGGACTCAC AGATTGATAA GCTAATGTTT
3251 AGAGAATTTA GATCGGAGAG AGTCGGTACG GCGCAGACTC AACATCAACC
3301 TCTTGCAAGC AACTAAAATG GCCTCGTCCT TGCTGTTTAT AACAGAAAAC
3351 AGACTTGTA AAAGCTTAGA TCATCAAGTG TTTTGGATTG GGGGCCTCCC
3401 AAAGGGATAT AAGAGGGGCA GGCCACTCTT AAGAAGAATG CGAGCTTTCT
3451 ACATTGGGAC TAGCATAAGA TCAAAGCCAA TCAAGATGGA GCACAGTAAC
3501 AGAAAACGTC GGTTCCTGTG GGAGAACAGA AGGGGAAAGG GTCTTAACTG
3551 GGAAAGGGCT CTGTGTGTA ACACCTCAGT TGTGTTCTCC TGACACCAGG
3601 AAAAGAGAGG GATCAGCTTC AATAACTAGA AAATTCTGGC TGTTTAATGG
3651 ACTCTTTGGT GGCTCTTTA AGGCAAAGCA GAGAAAGCAA ATTATGTATT
3701 AAGTGATTTT TGCATTTTAA AAACCTGACG TGCTGTATTG TACTAAATTA
3751 AGTGTAATCT ATTAAGGCAA GGTATACACA ATTTGCTTTG AAACCTACTA
3801 TGTTTATTCT ATTATAAAGT GTATTCAGGT GCAACACAGA GACTGCTTTC
3851 GGTGACATTA ATGAAGAAAA TTTCTCATGC CAGGCTTTAT TATAGAATCT
3901 TCAGCTAAAA TCCTAACTTT CTCCTTATTT CTTGGCACTT GTATACAAAGT
3951 GGTGTTGCCT CTTAGGGCAG GCATGAGCTA TTCTTTTCTG TAAAAATTTT
4001 TGAATCTATA GGCTGTGGGT TTCATTTTTG AAAAGTATTT TGTCTGGATG
4051 TCTTTCAAAC TAGCTTCAGA TATTATTTAA TACTATGTAA CTGGGTCCCC
4101 TATGGCTCAA TCAATATTGC TTATTTTTCT TCTGTAGTGG ATGTGAAATT
4151 TCCTTTAGTT GGATAAGATA CACTGTAATA ATTTTAATGC TAATTAATGA
4201 TATTTCATAC TGTGCAATGA ACAGATAATT TAACACTGTA TTTTGAAATG
4251 TTTTTTTCTT CCTGTCACCG CAGTGTGTGG TATTGCATAA TGTGAATACC
4301 TGTAAAAATA TAAATTACTT AAAAATAAAA ATATGACCAA TTGGTATCAG

-19/26-

Figure 15: SEQ ID NO. 7

Length: 50 bp

1 GGAGAGAGGCAGCACTGTAAACTGAAGTCAAATAAATTCAGCTCTTAATG


-20/26-

**Fig. 16: Alignment of SEQ ID NO. 7
with human MAGUIN-1 cDNA**

Length: 50 bp

```
      .       .       .       .  
1  GGAGAGAGGCAGCACTGTAAACTGAAGTCAAATAAATTCAGCTCTTAATG 50  
  ||| | | | | | | | | | | | | | | | | | | | | | | | | | | |  
5693 GGAGAGAGGCAGCACTGTAAACTGAAGTCAAATAAATTCAGCTCTTAATG 5742  
      .       .       .       .
```

Fig. 17: Schematic alignment of SEQ ID NO. 5, SEQ ID NO. 6 and SEQ ID NO. 7 with Genome Database EST-cluster



-22/26-

**Fig. 18: Images of the human cerebral cortex
labeled with anti-Maguin-1 antiserum
and with DAPI**




Table 1:

sample	Δ (fold) (frontal / temporal cortex)
patient P012	2.46
patient P016	2.78
patient P010	4.14
patient P011	2.20
patient P014	1.48
patient P017	1.42
patient P019	1.68
control C011	1.28
control C012	1.29
control C014	0.30
control C005	1.36
control C008	1.15

-24/26-**Table 2:**

sample	Δ (fold) (frontal cortex / hippocampus)
patient P012	1.37
patient P016	3.07
patient P010	2.99
patient P011	2.28
patient P014	1.21
patient P019	1.48
control C005	1.74
control C008	0.39
control C004	0.87

-25/26-**Table 3:**

sample	Δ (fold) (frontal / temporal cortex)
patient P012	2.68
patient P016	2.72
patient P010	11.73
patient P011	2.44
patient P014	1.77
patient P017	3.43
patient P019	4.02
control C011	1.42
control C012	1.22
control C014	0.30
control C005	0.92
control C008	0.81

-26/26-**Table 4:**

sample	Δ (fold) (frontal cortex / hippocampus)
patient P012	1.57
patient P016	4.38
patient P010	9.08
patient P011	4.53
patient P014	0.72
patient P019	1.37
control C005	1.84
control C008	0.46
control C004	1.69

SEQUENCE LISTING

<110> Evotec NeuroSciences GmbH

<120> DIAGNOSTIC AND THERAPEUTIC USE OF HUMAN MAGUIN PROTEINS
AND NUCLEIC ACIDS FOR NEURODEGENERATIVE DISEASES

<130> 030640wo ME/BM

<140> 02006353.3

<141> 2002-03-21

<160> 24

<170> PatentIn Ver. 2.1

<210> 1

<211> 1034

<212> PRT

<213> Homo sapiens

<400> 1

Met Ala Leu Ile Met Glu Pro Val Ser Lys Trp Ser Pro Ser Gln Val
1 5 10 15

Val Asp Trp Met Lys Gly Leu Asp Asp Cys Leu Gln Gln Tyr Ile Lys
20 25 30

Asn Phe Glu Arg Glu Lys Ile Ser Gly Asp Gln Leu Leu Arg Ile Thr
35 40 45

His Gln Glu Leu Glu Asp Leu Gly Val Ser Arg Ile Gly His Gln Glu
50 55 60

Leu Ile Leu Glu Ala Val Asp Leu Leu Cys Ala Leu Asn Tyr Gly Leu
65 70 75 80

Glu Thr Glu Asn Leu Lys Thr Leu Ser His Lys Leu Asn Ala Ser Ala
85 90 95

Lys Asn Leu Gln Asn Phe Ile Thr Gly Arg Arg Arg Ser Gly His Tyr
100 105 110

Asp Gly Arg Thr Ser Arg Lys Leu Pro Asn Asp Phe Leu Thr Ser Val
115 120 125

Val Asp Leu Ile Gly Ala Ala Lys Ser Leu Leu Ala Trp Leu Asp Arg
130 135 140

Ser Pro Phe Ala Ala Val Thr Asp Tyr Ser Val Thr Arg Asn Asn Val
145 150 155 160

Ile Gln Leu Cys Leu Glu Leu Thr Thr Ile Val Gln Gln Asp Cys Thr
165 170 175

Val Tyr Glu Thr Glu Asn Lys Ile Leu His Val Cys Lys Thr Leu Ser
180 185 190

Gly Val Cys Asp His Ile Ile Ser Leu Ser Ser Asp Pro Leu Val Ser
195 200 205

Gln Ser Ala His Leu Glu Val Ile Gln Leu Ala Asn Ile Lys Pro Ser
210 215 220

Glu Gly Leu Gly Met Tyr Ile Lys Ser Thr Tyr Asp Gly Leu His Val
225 230 235 240

Ile Thr Gly Thr Thr Glu Asn Ser Pro Ala Asp Arg Cys Lys Lys Ile
245 250 255

His Ala Gly Asp Glu Val Ile Gln Val Asn His Gln Thr Val Val Gly
260 265 270

Trp Gln Leu Lys Asn Leu Val Asn Ala Leu Arg Glu Asp Pro Ser Gly
275 280 285

Val Ile Leu Thr Leu Lys Lys Arg Pro Gln Ser Met Leu Thr Ser Ala
290 295 300

Pro Ala Leu Leu Lys Asn Met Arg Trp Lys Pro Leu Ala Leu Gln Pro
305 310 315 320

Leu Ile Pro Arg Ser Pro Thr Ser Ser Val Ala Thr Pro Ser Ser Thr
325 330 335

Ile Ser Thr Pro Thr Lys Arg Asp Ser Ser Ala Leu Gln Asp Leu Tyr
340 345 350

Ile Pro Pro Pro Pro Ala Glu Pro Tyr Ile Pro Arg Asp Glu Lys Gly
355 360 365

Asn Leu Pro Cys Glu Asp Leu Arg Gly His Met Val Gly Lys Pro Val
370 375 380

His Lys Gly Ser Glu Ser Pro Asn Ser Phe Leu Asp Gln Glu Tyr Arg
385 390 395 400

Lys Arg Phe Asn Ile Val Glu Glu Asp Thr Val Leu Tyr Cys Tyr Glu
405 410 415

Tyr Glu Lys Gly Arg Ser Ser Ser Gln Gly Arg Arg Glu Ser Thr Pro
420 425 430

Thr Tyr Gly Lys Leu Arg Pro Ile Ser Met Pro Val Glu Tyr Asn Trp
435 440 445

Val Gly Asp Tyr Glu Asp Pro Asn Lys Met Lys Arg Asp Ser Arg Arg
450 455 460

Glu Asn Ser Leu Leu Arg Tyr Met Ser Asn Glu Lys Ile Ala Gln Glu
465 470 475 480

Glu Tyr Met Phe Gln Arg Asn Ser Lys Lys Asp Thr Gly Lys Lys Ser
485 490 495

Lys Lys Lys Gly Asp Lys Ser Asn Ser Pro Thr His Tyr Ser Leu Leu
500 505 510

Pro Ser Leu Gln Met Asp Ala Leu Arg Gln Asp Ile Met Gly Thr Pro
515 520 525

Val Pro Glu Thr Thr Leu Tyr His Thr Phe Gln Gln Ser Ser Leu Gln
530 535 540

His Lys Ser Lys Lys Lys Asn Lys Gly Pro Ile Ala Gly Lys Ser Lys
545 550 555 560

Arg Arg Ile Ser Cys Lys Asp Leu Gly Arg Gly Asp Cys Glu Gly Trp
565 570 575

Leu Trp Lys Lys Lys Asp Ala Lys Ser Tyr Phe Ser Gln Lys Trp Lys
580 585 590

Lys Tyr Trp Phe Val Leu Lys Asp Ala Ser Leu Tyr Trp Tyr Ile Asn
595 600 605

Glu Glu Asp Glu Lys Ala Glu Gly Phe Ile Ser Leu Pro Glu Phe Lys
610 615 620

Ile Asp Arg Ala Ser Glu Cys Arg Lys Lys Tyr Ala Phe Lys Ala Cys
625 630 635 640

His Pro Lys Ile Lys Ser Phe Tyr Phe Ala Ala Glu His Leu Asp Asp
645 650 655

Met Asn Arg Trp Leu Asn Arg Ile Asn Met Leu Thr Ala Gly Tyr Ala
660 665 670

Glu Arg Glu Arg Ile Lys Gln Glu Gln Asp Tyr Trp Ser Glu Ser Asp
675 680 685

Lys Glu Glu Ala Asp Thr Pro Ser Thr Pro Lys Gln Asp Ser Pro Pro
690 695 700

Pro Pro Tyr Asp Thr Tyr Pro Arg Pro Pro Ser Met Ser Cys Ala Ser
705 710 715 720

Pro Tyr Val Glu Ala Lys His Ser Arg Leu Ser Ser Thr Glu Thr Ser
725 730 735

Gln Ser Gln Ser Ser His Glu Glu Phe Arg Gln Glu Val Thr Gly Ser
740 745 750

Ser Ala Val Ser Pro Ile Arg Lys Thr Ala Ser Gln Arg Arg Ser Trp
755 760 765

Gln Asp Leu Ile Glu Thr Pro Leu Thr Ser Ser Gly Leu His Tyr Leu
770 775 780

Gln Thr Leu Pro Leu Glu Asp Ser Val Phe Ser Asp Ser Ala Ala Ile
785 790 795 800

Ser Pro Glu His Arg Arg Gln Ser Thr Leu Pro Thr Gln Lys Cys His
805 810 815

Leu Gln Asp His Tyr Gly Pro Tyr Pro Leu Ala Glu Ser Glu Arg Met
820 825 830

Gln Val Leu Asn Gly Asn Gly Gly Lys Pro Arg Ser Phe Thr Leu Pro
835 840 845

Arg Asp Ser Gly Phe Asn His Cys Cys Leu Asn Ala Pro Val Ser Ala
850 855 860

Cys Asp Pro Gln Asp Asp Val Gln Pro Pro Glu Val Glu Glu Glu Glu
865 870 875 880

Glu Glu Glu Glu Glu Glu Gly Glu Ala Ala Gly Glu Asn Ile Gly Glu
885 890 895

Lys Ser Glu Ser Arg Glu Glu Lys Leu Gly Asp Ser Leu Gln Asp Leu
900 905 910

Tyr Arg Ala Leu Glu Gln Ala Ser Leu Ser Pro Leu Gly Glu His Arg
915 920 925

Ile Ser Thr Lys Met Glu Tyr Lys Leu Ser Phe Ile Lys Arg Cys Asn
930 935 940

Asp Pro Val Met Asn Glu Lys Leu His Arg Leu Arg Ile Leu Lys Ser
945 950 955 960

Thr Leu Lys Ala Arg Glu Gly Glu Val Ala Ile Ile Asp Lys Val Leu
965 970 975

Asp Asn Pro Asp Leu Thr Ser Lys Glu Phe Gln Gln Trp Lys Gln Met
980 985 990

Tyr Leu Asp Leu Phe Leu Asp Ile Cys Gln Asn Thr Thr Ser Asn Asp
995 1000 1005

Pro Leu Ser Ile Ser Ser Glu Val Asp Val Ile Thr Ser Ser Leu Ala
1010 1015 1020

His Thr His Ser Tyr Ile Glu Thr His Val
1025 1030

<210> 2
 <211> 948
 <212> PRT
 <213> Homo sapiens

<400> 2
 Phe Ile Gly Arg Glu Ser Glu Gln Ile Asp Asn Ala Met Ile Asn Ala
 1 5 10 15
 Cys Ile Asp Ser Glu Gln Glu Asn Cys Glu Phe His Met Ala Asn Met
 20 25 30
 Ala Gly Ile Asn Pro Arg Thr Glu Ile Asn Leu Glu Asn Gly Thr His
 35 40 45
 Ala Ala Met Ala Leu Ile Met Glu Pro Val Ser Lys Trp Ser Pro Ser
 50 55 60
 Gln Val Val Asp Trp Met Lys Gly Leu Asp Asp Cys Leu Gln Gln Tyr
 65 70 75 80
 Ile Lys Asn Phe Glu Arg Glu Lys Ile Ser Gly Asp Gln Leu Leu Arg
 85 90 95
 Ile Thr His Gln Glu Leu Glu Asp Leu Gly Val Ser Arg Ile Gly His
 100 105 110
 Gln Glu Leu Ile Leu Glu Ala Val Asp Leu Leu Cys Ala Leu Asn Tyr
 115 120 125
 Gly Leu Glu Thr Glu Asn Leu Lys Thr Leu Ser His Lys Leu Asn Ala
 130 135 140
 Ser Ala Lys Asn Leu Gln Asn Phe Ile Thr Gly Arg Arg Arg Ser Gly
 145 150 155 160
 His Tyr Asp Gly Arg Thr Ser Arg Lys Leu Pro Asn Asp Phe Leu Thr
 165 170 175
 Ser Val Val Asp Leu Ile Gly Ala Ala Lys Ser Leu Leu Ala Trp Leu
 180 185 190
 Asp Arg Ser Pro Phe Ala Ala Val Thr Asp Tyr Ser Val Thr Arg Asn
 195 200 205
 Asn Val Ile Gln Leu Cys Leu Glu Leu Thr Thr Ile Val Gln Gln Asp
 210 215 220
 Cys Thr Val Tyr Glu Thr Glu Asn Lys Ile Leu His Val Cys Lys Thr
 225 230 235 240
 Leu Ser Gly Val Cys Asp His Ile Ile Ser Leu Ser Ser Asp Pro Leu

245 250 255

Val Ser Gln Ser Ala His Leu Glu Val Ile Gln Leu Ala Asn Ile Lys
260 265 270

Pro Ser Glu Gly Leu Gly Met Tyr Ile Lys Ser Thr Tyr Asp Gly Leu
275 280 285

His Val Ile Thr Gly Thr Thr Glu Asn Ser Pro Ala Asp Arg Cys Lys
290 295 300

Lys Ile His Ala Gly Asp Glu Val Ile Gln Val Asn His Gln Thr Val
305 310 315 320

Val Gly Trp Gln Leu Lys Asn Leu Val Asn Ala Leu Arg Glu Asp Pro
325 330 335

Ser Gly Val Ile Leu Thr Leu Lys Lys Arg Pro Gln Ser Met Leu Thr
340 345 350

Ser Ala Pro Ala Leu Leu Lys Asn Met Arg Trp Lys Pro Leu Ala Leu
355 360 365

Gln Pro Leu Ile Pro Arg Ser Pro Thr Ser Ser Val Ala Thr Pro Ser
370 375 380

Ser Thr Ile Ser Thr Pro Thr Lys Arg Asp Ser Ser Ala Leu Gln Asp
385 390 395 400

Leu Tyr Ile Pro Pro Pro Pro Ala Glu Pro Tyr Ile Pro Arg Asp Glu
405 410 415

Lys Gly Asn Leu Pro Cys Glu Asp Leu Arg Gly His Met Val Gly Lys
420 425 430

Pro Val His Lys Gly Ser Glu Ser Pro Asn Ser Phe Leu Asp Gln Glu
435 440 445

Tyr Arg Lys Arg Phe Asn Ile Val Glu Glu Asp Thr Val Leu Tyr Cys
450 455 460

Tyr Glu Tyr Glu Lys Gly Arg Ser Ser Ser Gln Gly Arg Arg Glu Ser
465 470 475 480

Thr Pro Thr Tyr Gly Lys Leu Arg Pro Ile Ser Met Pro Val Glu Tyr
485 490 495

Asn Trp Val Gly Asp Tyr Glu Asp Pro Asn Lys Met Lys Arg Asp Ser
500 505 510

Arg Arg Glu Asn Ser Leu Leu Arg Tyr Met Ser Asn Glu Lys Ile Ala
515 520 525

Gln Glu Glu Tyr Met Phe Gln Arg Asn Ser Lys Lys Asp Thr Gly Lys

530	535	540
Lys Ser Lys Lys Lys Gly Asp Lys Ser Asn Ser Pro Thr His Tyr Ser		
545	550	555 560
Leu Leu Pro Ser Leu Gln Met Asp Ala Leu Arg Gln Asp Ile Met Gly		
	565	570 575
Thr Pro Val Pro Glu Thr Thr Leu Tyr His Thr Phe Gln Gln Ser Ser		
	580	585 590
Leu Gln His Lys Ser Lys Lys Lys Asn Lys Gly Pro Ile Ala Gly Lys		
	595	600 605
Ser Lys Arg Arg Ile Ser Cys Lys Asp Leu Gly Arg Gly Asp Cys Glu		
	610	615 620
Gly Trp Leu Trp Lys Lys Lys Asp Ala Lys Ser Tyr Phe Ser Gln Lys		
	625	630 635 640
Trp Lys Lys Tyr Trp Phe Val Leu Lys Asp Ala Ser Leu Tyr Trp Tyr		
	645	650 655
Ile Asn Glu Glu Asp Glu Lys Ala Glu Gly Phe Ile Ser Leu Pro Glu		
	660	665 670
Phe Lys Ile Asp Arg Ala Ser Glu Cys Arg Lys Lys Tyr Ala Phe Lys		
	675	680 685
Ala Cys His Pro Lys Ile Lys Ser Phe Tyr Phe Ala Ala Glu His Leu		
	690	695 700
Asp Asp Met Asn Arg Trp Leu Asn Arg Ile Asn Met Leu Thr Ala Gly		
	705	710 715 720
Tyr Ala Glu Arg Glu Arg Ile Lys Gln Glu Gln Asp Tyr Trp Ser Glu		
	725	730 735
Ser Asp Lys Glu Glu Ala Asp Thr Pro Ser Thr Pro Lys Gln Asp Ser		
	740	745 750
Pro Pro Pro Pro Tyr Asp Thr Tyr Pro Arg Pro Pro Ser Met Ser Cys		
	755	760 765
Ala Ser Pro Tyr Val Glu Ala Lys His Ser Arg Leu Ser Ser Thr Glu		
	770	775 780
Thr Ser Gln Ser Gln Ser Ser His Glu Glu Phe Arg Gln Glu Val Thr		
	785	790 795 800
Gly Ser Ser Ala Val Ser Pro Ile Arg Lys Thr Ala Ser Gln Arg Arg		
	805	810 815
Ser Trp Gln Asp Leu Ile Glu Thr Pro Leu Thr Ser Ser Gly Leu His		

820 825 830
 Tyr Leu Gln Thr Leu Pro Leu Glu Asp Ser Val Phe Ser Asp Ser Ala
 835 840 845
 Ala Ile Ser Pro Glu His Arg Arg Gln Ser Thr Leu Pro Thr Gln Lys
 850 855 860
 Cys His Leu Gln Asp His Tyr Gly Pro Tyr Pro Leu Ala Glu Ser Glu
 865 870 875 880
 Arg Met Gln Val Leu Asn Gly Asn Gly Gly Lys Pro Arg Ser Phe Thr
 885 890 895
 Leu Pro Arg Asp Ser Gly Phe Asn His Cys Cys Leu Asn Ala Pro Val
 900 905 910
 Ser Ala Cys Asp Pro Gln Asp Asp Val Gln Pro Pro Glu Val Glu Glu
 915 920 925
 Glu Glu Glu Glu Glu Glu Glu Glu Gly Glu Ala Ala Gly Glu Asn Ile
 930 935 940
 Gly Glu Lys Ser
 945

<210> 3
 <211> 3105
 <212> DNA
 <213> Homo sapiens

<400> 3
 atggctctga taatggaacc ggtgagcaaa tgggtctccga gtcaagtagt ggactggatg 60
 aaaggtcttg atgactgttt gcagcagtat attaagaact ttgagagggga gaagatcagt 120
 ggggaccagc tgctgcgcat tacacatcag gagctagaag atctgggggt cagccgcatt 180
 ggccatcagg aactgatctt ggaagcagtt gaccttctgt gtgcattgaa ttatggcttg 240
 gaaacagaaa atctaaaaac cctttctcac aagttgaatg catctgccaa aaatctgcag 300
 aattttataa caggaaggag aaggagtggc cattatgatg ggaggaccag cggaaaattg 360
 ccaaacgact ttctgacctc agttgtggat ctgattggag cagccaagag tctgcttgcc 420
 tggttggaca ggtcaccatt tgctgctgtg acagactatt cagttacaag aaataatgtc 480
 atacaactct gcctggagtt aacaacaatt gtgcaacagg attgtactgt atatgaaaca 540
 gagaataaaa ttcttcacgt gtgtaaaact ctttctggag tctgtgacca catcatatcc 600
 ctgtcgtcag atcctctggg ttacacagtct gctcacctgg aagtgattca actggcaaac 660
 attaaaccaa gcgaagggct gggatgtat attaaatcta catatgatgg cctccatgta 720
 attactggaa ccacagaaaa ttcacctgca gatcgggtgca agaaaatcca tgctggcgat 780
 gaagtgattc aagttaatca tcagactgtg gtgggggtggc agttgaaaaa tttgggtgaat 840
 gcactacgag aggaccagag tgggtgttat ttaactttga aaaagcgacc tcagagcatg 900
 cttacctcag caccagcttt actgaaaaat atgagatgga agcccttgc tctgcagcct 960
 cttataccta gaagtccac aagcagcgtt gccacgcctt ccagcaccat cagtacacc 1020
 accaaaagag acagttctgc cctccaggat ctctacattc cccctcctcc tgcagaacca 1080
 tatattccca gggatgaaaa aggaaacctt ccttgtgaag acctcagagg acatattggtg 1140
 ggcaagccag tgcataaggg atctgaatca ccaaattcat ttctggatca ggaatatcga 1200
 aagagattta atattgtcga agaagatact gtcttatatt gctatgaata tgaaaaagga 1260

```

agatcaagta gtcaaggaag acgagaaagc accccaactt atggcaagct acgacctata 1320
tctatgccag tggaatataa ttgggtgggg gactatgaag atccaaataa gatgaagaga 1380
gatagtagaa gagaaaactc tctacttcgg tatatgagca atgaaaagat tgctcaagaa 1440
gaatacatgt ttcagagaaa cagcaaaaag gacacaggga agaagtcaaa aaagaagggt 1500
gataagagta atagcccaac tcactattca ttgctaccta gtttacaat ggatgcactg 1560
agacaagaca tcatgggcac tcctgtgcc aagaccacac tataccatac atttcagcag 1620
tcctcactgc agcacaaatc aaagaagaaa aacaaaggct ctatagcagg caagagcaaa 1680
agacgaattt cttgcaaaga tcttgccgt ggtgactgtg agggctggct ttggaaaaag 1740
aaagatgcga agagttactt ttcacagaaa tggaaaaaat attggtttgt cctaaaggat 1800
gcatcccttt attggtatat taatgaggag gatgaaaaag cagaaggatt cattagcctg 1860
cctgaattta aaattgatag agccagtga tgcgcagaaa aatatgcatt caaagcctgt 1920
catcctaaaa tcaaaagctt ttattttgt gctgaacatc ttgatgatat gaacagggtg 1980
cttaacagaa ttaatatgct gactgcagga tatgcagaaa gagagaggat taagcaggaa 2040
caagattact ggagtgcag tgacaaggaa gaagcagata ctccatcaac accaaaacaa 2100
gatagccctc caccgccata tgatacata ccacgacctc cctcgatgag ttgcccag 2160
ccttatgttg aagcaaaaaca tagccgactt tcctccacgg agacttctca gtctcagtct 2220
tctcatgagg agtttcgcca ggaagtaact gggagcagtg cagtgtctcc cattcgcaag 2280
acagccagtc agcgcgcgtc ctggcaggat ttaattgaga cgccactgac aagttcaggc 2340
ttacactatc ttcagactct gccctggag gattctgtct tctctgactc cgcgccatc 2400
tccccagagc acaggcggca gtctaccctg ccaactcaga aatgccacct gcaggatcac 2460
tatgggcat accccttagc tgagagtgc aggatgcaag tgctaaatgg aaatgggggc 2520
aagcctcgaa gttttactct gcctcgagat agcgggttca accattgctg tctgaatgct 2580
ccagttagtg cctgtgacct acaggatgac gtgcaacccc cagagggtga ggaagaggag 2640
gaggaggagg aggaggaagg ggaggcagca ggggaaaaa taggagaaaa aagtgaagc 2700
agagaagaaa agttaggaga ctcatgcaa gatttataca gggcactgga gcaggccagt 2760
ctgtcaccac taggagaaca tcgtatttca accaagatgg aatacaagct atcattata 2820
aaaagatgta atgatcctgt aatgaatgaa aaactacacc ggctgagaat tctcaaaagc 2880
actttaaagg ccagagaagg ggaagtagcc attatcgata aagtcctaga caatccagac 2940
ttgacatcta aagaattcca acaatggaag cagatgtacc tcgacctttt cttggatata 3000
tgtcaaaaata ccacctcaa tgacctactg agtatttctt ctgaagtaga tgtaatcact 3060
tcctctctag cacacactca ttcatacatt gaaacgcag tctaa 3105

```

<210> 4

<211> 2683

<212> DNA

<213> Homo sapiens

<400> 4

```

atggctctga taatggaacc ggtgagcaaa tggctctccga gtcaagtagt ggactggatg 60
aaaggtcttg atgactgttt gcagcagtat attaagaact ttgagaggga gaagatcagt 120
ggggaccagc tgctgcgcat tacacatcag gagctagaag atctgggggt cagccgcatt 180
ggccatcagg aactgatctt ggaagcagtt gaccttctgt gtgcattgaa ttatggcttg 240
gaaacagaaa atctaaaaac cctttctcac aagttgaatg catctgcaa aaatctgcag 300
aattttataa caggaaggag aaggagtggc cattatgatg ggaggaccag ccgaaaattg 360
ccaaacgact ttctgacctc agttgtggat ctgattggag cagccaagag tctgcttgcc 420
tggttgagca ggtcaccatt tgctgctgtg acagactatt cagttacaag aaataatgtc 480
atacaactct gcctggagtt aacaacaatt gtgcaacagg attgtactgt atatgaaaca 540
gagaataaaa ttcttcacgt gtgtaaaact ctttctggag tctgtgacca catcatatcc 600
ctgtcgtcag atcctctggt ttcacagtct gctcacctgg aagtgattca gctggcaaac 660
attaaccaca gcgaagggtt gggtatgtat attaaatcta catatgatgg cctccatgta 720
attactggaa ccacagaaaa ttcacctgca gatcgggtgca agaaaatcca tgctggcgat 780
gaagtgatto aagttaatca tcagactgtg gtgggggtggc agttgaaaaa tttggtgaat 840
gcactacgag aggacctgag tgggtgttatc ttaactttga aaaagcgacc tcagagcatg 900

```

```
cttacctcag caccagcttt actgaaaaat atgagatgga agcccccttg tctgcagcct 960
cttataccta gaagtcccac aagcagcggt gccacgcctt ccagcaccat cagtacaccc 1020
accaaaagag acagttctgc cctccaggat ctctacattc cccctcctcc tgcagaacca 1080
tatattccca gggatgaaaa aggaaacctt ccttgtgaag acctcagagg acatatggtg 1140
ggcaagccag tgcataaggg atctgaatca ccaaattcat ttctggatca ggaatatcga 1200
aagagattta atattgtcga agaagatact gtcttatatt gctatgaata tgaaaaagga 1260
agatcaagta gtcaagggaag acgagaaagc accccaactt atggcaagct acgacctata 1320
tctatgccag tggaatataa ttgggtgggg gactatgaag atccaaataa gatgaagaga 1380
gatagtagaa gagaaaactc tctacttcgg tatatgagca atgaaaagat tgctcaagaa 1440
gaatacatgt ttcagagaaa cagcaaaaag gacacaggga agaagtcaaa aaagaagggt 1500
gataagagta atagcccaac tcactattca ttgctaccta gtttacaat ggatgcactg 1560
agacaagaca tcatgggcac tcctgtgcca gagaccacac tataccatac atttcagcag 1620
tcctcactgc agcacaatc aaagaagaaa aacaaaggct ctatagcagg caagagcaaa 1680
agacgaattt cttgcaaaga tcttggccgt ggtgactgtg agggctggct ttggaaaaag 1740
aaagatgcga agagttactt ttcacagaaa tggaaaaaat attggtttgt cctaaaggat 1800
gcataccctt attggtatat taatgaggag gatgaaaaag cagaaggatt cattagcctg 1860
cctgaattta aaattgatag agccagtga tgccgcaaaa aatatgcatt caaagcctgt 1920
catcctaaaa tcaaaagctt ttattttgc tctgaacatc ttgatgatat gaacagggtg 1980
cttaacagaa ttaatatgct gactgcagga tatgcagaaa gagagaggat taagcaggaa 2040
caagattact ggagtgagag tgacaaggaa gaagcagata ctccatcaac accaaaacaa 2100
gatagccctc ccccccata tgatacatc ccacgacctc cctcgatgag ttgcgccagt 2160
ccttatgtgg aagcaaaaca tagccgactt tcctccacgg agacttctca gtctcagtct 2220
tctcatgagg agtttcgcca ggaagtaact gggagcagtg cagtgtctcc cattcgcaag 2280
acagccagtc agcgcgcctc ctggcaggat ttaattgaga cgccactgac aagttcaggc 2340
ttacactatc ttcagactct gcccctggag gattctgtct tctctgactc cgcggccatc 2400
tccccagagc acaggcggca gtctaccctg ccaactcaga aatgccacct gcaggatcac 2460
tatgggccat accccttagc tgagagttag aggatgcaag tgctaaatgg aaatgggggc 2520
aagcctcgaa gttttactct gcctcgagat agcgggttca accattgctg tctgaatgct 2580
ccagtttagt cctgtgacct acaggatgac gtgcaacccc cagagggtga ggaagaggag 2640
gaggaggagg aggaggaagg ggaggcagca ggggaaaaca tag 2683
```

<210> 5

<211> 5749

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA of human
Maguin-1

<400> 5

```
cgggcagcta gtcgtgctcg gggcttcact cccgcgcgtg aggcgagcgg gcaagttggc 60
tgagggcggt cggcagaggc tgcttccctc ggcgacgcga cccctcagca actcaagcta 120
tgaactgaag ctccctaggg acggagaccg gagcggagcg gcggaggcag cagcagcagc 180
agcagcagca gcagcagcag cagccgcgcg cgccgcgcgc ttagcgggaa ctgagcagac 240
ccggcgcgga gccacgactc ctgcacgttt acctccctgt cgcggttctt cgcggcggtt 300
ggctaaaaga cgttacagcc gcgagaccgc acacacaaaa gccgctttct ccgcgcgcgc 360
cgcccaggga ggctgcggcc agcaagggac cccacctgag agcagctcgg gctgctgagt 420
tcgttttgtg tctgagctct gcgctctgca cggaaccgac cccgtaccca tggctctgat 480
aatggaaccg gtgagcaaat ggtctccgag tcaagtagtg gactggatga aaggtcttga 540
tgactgtttg cagcagtata ttaagaactt tgagagggag aagatcagtg gggaccagct 600
gctgcgcatt acacatcagg agctagaaga tctgggggtc agccgcattg gccatcagga 660
actgatcttg gaagcagttg accttctgtg tgcattgaat tatggcttgg aaacagaaaa 720
```

tctaaaaacc	ctttctcaca	agttgaatgc	atctgccaaa	aatctgcaga	attttataac	780
aggaaggaga	aggagtggcc	attatgatgg	gaggaccagc	cgaaaattgc	caaacgactt	840
tctgacctca	gttgtggatc	tgattggagc	agccaagagt	ctgcttgcc	ggttggacag	900
gtcaccat	gctgctgtga	cagactattc	agttacaaga	aataatgtca	tacaactctg	960
cctggagtta	acaacaattg	tgcaacagga	ttgtactgta	tatgaaacag	agaataaaaat	1020
tcttcacgtg	tgtaaaactc	tttctggagt	ctgtgaccac	atcatatccc	tgctcgtcaga	1080
tcctctgggt	tcacagtctg	ctcacctgga	agtgattcaa	ctggcaaaca	ttaaaccaag	1140
cgaagggctg	ggtatgtata	ttaaatctac	atatgatggc	ctccatgtaa	ttactggaac	1200
cacagaaaat	tcacctgcag	atcggtgcaa	gaaaatccat	gctggcgatg	aagtgattca	1260
agttaatcat	cagactgtgg	tggggtggca	gttgaaaaat	ttggtgaatg	cactacgaga	1320
ggacccgagt	ggtgttatct	taactttgaa	aaagcgacct	cagagcatgc	ttacctcagc	1380
accagcttta	ctgaaaaata	tgagatggaa	gcccttgct	ctgcagcctc	ttatacctag	1440
aagtccca	agcagcgttg	ccacgccttc	cagcaccatc	agtacacca	ccaaaagaga	1500
cagttctgcc	ctccaggatc	tctacattcc	ccctcctcct	gcagaaccat	atattcccag	1560
ggatgaaaaa	ggaaaccttc	cttgtgaaga	cctcagagga	catatgggtg	gcaagccagt	1620
gcataaggga	tctgaatcac	caaattcatt	tctggatcag	gaatatcgaa	agagatttaa	1680
tattgtcgaa	gaagatactg	tcttatattg	ctatgaatat	gaaaaaggaa	gatcaagtag	1740
tcaaggaaga	cgagaaagca	ccccaaacta	tggcaagcta	cgacctatat	ctatgccagt	1800
ggaatataat	tgggtggggg	actatgaaga	tccaaataag	atgaagagag	atagtagaag	1860
agaaaactct	ctacttcggt	atatgagcaa	tgaâaagatt	gctcaagaag	aatacatgtt	1920
tcagagaaac	agcaaaaagg	acacagggaa	gaagtcaaaa	aagaaggggtg	ataagagtaa	1980
tagcccaact	cactattcat	tgctacctag	tttacaaatg	gatgcactga	gacaagacat	2040
catgggcact	cctgtgccag	agaccacact	ataccataca	tttcagcagt	cctcactgca	2100
gcacaaatca	aagaagaaaa	acaaaggtcc	tatagcaggg	aagagcaaaa	gacgaatttc	2160
ttgcaaagat	cttggccgtg	gtgactgtga	gggctggctt	tggaaaaaga	aagatgcgaa	2220
gagttacttt	tcacagaaat	ggaaaaaata	ttggtttgtc	ctaaaggatg	catcccttta	2280
ttggtatatt	aatgaggagg	atgaaaaagc	agaaggattc	attagcctgc	ctgaatttaa	2340
aattgataga	gccagtgaat	gccgcaaaa	atatgcattc	aaagcctgtc	atcctaaaat	2400
caaaagcttt	tattttgtcg	ctgaacatct	tgatgatatg	aacagggtgg	ttaacagaat	2460
taatgtgctg	actgcaggat	atgcagaaag	agagaggatt	aagcaggaac	aagattactg	2520
gagtgaagat	gacaaggaag	aagcagatac	tccatcaaca	ccaaaacaag	atagccctcc	2580
accccatat	gatacatacc	cacgacctcc	ctcgatgagt	tgcgccagtc	cttatgtgga	2640
agcaaaacat	agccgacttt	cctccacgga	gacttctcag	tctcagtctt	ctcatgagga	2700
gtttcgccag	gaagtaactg	ggagcagtg	agtgtctccc	attcgcaaga	cagccagtca	2760
gcgccgtcc	tggcaggatt	taattgagac	gccactgaca	agttcaggct	tacactatct	2820
tcagactctg	cccctggagg	attctgtctt	ctctgactcc	gcggccatct	ccccagagca	2880
caggcggcag	tctaccctgc	caactcagaa	atgccacctg	caggatcact	atgggccata	2940
ccccttagct	gagagtgaga	ggatgcaagt	gctaaatgga	aatgggggca	agcctcgaag	3000
ttttactctg	cctcgagata	gcgggttcaa	ccattgctgt	ctgaatgctc	cagttagtgc	3060
ctgtgaccca	caggatgacg	tgcaaccccc	agaggtggag	gaagaggagg	aggaggagga	3120
ggaggaaggg	gaggcagcag	gggaaaacat	aggagaaaaa	agtgaagca	gagaagaaaa	3180
gttaggagac	tcattgcaag	atttatacag	ggcactggag	caggccagtc	tgtcaccact	3240
aggagaacat	cgtatttcaa	ccaagatgga	atacaagcta	tcatttataa	aaagatgtaa	3300
tgatcctgta	atgaatgaaa	aactacaccg	gctgagaatt	ctcaaaagca	ctttaaaggc	3360
cagagaaggg	gaagtagcca	ttatcgataa	agtcctagac	aatccagact	tgacatctaa	3420
agaattccaa	caatggaagc	agatgtacct	cgaccttttc	ttggatatct	gtcaaaatac	3480
cacctcaaat	gaccactga	gtatttcttc	tgaagtagat	gtaatcactt	cctctctagc	3540
acacactcat	tcatacattg	aaacgcattg	ctaaatgtat	tctgccttca	gaccatctag	3600
tacctgctgg	tactctgaac	aagtatataa	ggtagttttt	atatcaatgt	gtggaacact	3660
tgacaagcta	tactttaatg	ttaccaaact	atatgaaaca	aaccatatat	ggtcacaata	3720
ccactatctt	taatgagcat	ttgtatat	tatatgcaac	agtgtctcag	ttatgtttac	3780
catgtgcaaa	atcaactgtc	tttaatgaact	taaaattaac	ttttgcaaac	aattctaaat	3840
acaggtgggtc	ttcaagtagt	aaaaccacaa	aaggcagttt	tctatctatg	gtcatctttt	3900
ctcccttttaa	gttaatttta	tataaacaag	acttcaaaag	taaatcacat	tttttcaggt	3960

```

gcagacatcc ttgtgggtgg gaaagaatth aaacctthtt tatatthatt aaaatgttct 4020
aagaatthttc ttaaaccattg caccaaagtht aatgctgtag thttatthtt gtgaaatgta 4080
gatgcgcata caagagctaa gcaaaataga agagcatcga cataagaaaa gttcagggtat 4140
ctaattattcg tcttaatatg ctattaactt gtgaaagcta agttaatgga aatattattc 4200
caaattctatg agaacacttg gtgtatcagg gcaaagctth gtaagatgtt thtgtaacta 4260
agacccaaat tgaagataga gctgctthtt thttcttggt taaatcttcc thttatthttg 4320
tagtgatgag atgctgattg tgtacagaag aatthtgagag gggatthttt aaaactgact 4380
taacacaccc agaaaggcag ctaacagcta tatatatata taaatthcag cccaaactca 4440
tgthttttaa ctccaactct taaaagacaa caaggtataa actgaaatga atcaactthc 4500
cacttagtht ccaatthttc cctagthccac taatthaaact taggtaatta tacttcagggt 4560
aggaagtac aatatgttht gthtcaggct gatgtgtgtt ataaaaaca acactgaaaa 4620
ataaaattgt actthccctt taaggagcaa gcagggtgat gtcattcaaa gagatgtcac 4680
attgaattat gagagaaaca atthtagaggt thttthctct gcttcatgaa thgttctata 4740
gagtgatga agtctaagga aaagthctct tcatatattt ccattthata gcgtcttgtt 4800
thtgaaagtg atcacagcat gaaaatgact gtgctgctth thagtgtctg gctgcataat 4860
gtacaagtca caatthgtg thttthttcag gaggagaaag ggaacctctt ttactattct 4920
atathctaaa atctacttht aatcagctth atactgttg ctgtacagct cagtgaatgt 4980
actthctatct ttaagagthc agatathgct cagtgaatat thttgctgta gaggagaaag 5040
taaaaactcc acagcgggga thttthttct tgctthtgaa accaccattg aatcactatc 5100
gthttgcaga cthttgcacaa ctgtacagga gagtggtctt tctacagcac atthttcagta 5160
atctatatt tagtcaaaat ggatgagaaa tcatgtatta atgtthgtat ggaatthttg 5220
gtccagtgt ataththttt cattthaaaa gaactctatt tgtaaaaaa thttatthact 5280
gcatggatat tgacgcacat taaatthgtg ggatthttgt tatgtaaaaa aaaaaaaaaa 5340
aaaaaaaaaac aaaaaacctt thgtctctaaa atgaagtgtg cthgttaaca ggtgtthtaga 5400
cttattgatg thtactagac caaatgtgt thgttactth aaaatathatg tacctgatgg 5460
atgtgtcatg thtacagtgg ccaggthgtg gcctgtaaac agcaagcagt tgacgggaag 5520
actagctctg thgtactaa gcagctthtt cthttgtaaa gtcagctctg thgtthttaaa 5580
tggtaaaaat taaactaatg aatthgacaa gactcgtggc tagcctagca tgaaagagac 5640
ctthtaacac tatataatat ctgtacattt tathgcatth gthttcaaatc taggagagag 5700
gcagcactgt aaactgaagt caaataaatt cagctcttaa tgaatctth 5749

```

<210> 6

<211> 4350

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA of human
Maguin-2

<400> 6

```

gtgctcgggg cttcactccc gcgcgtgagg cgagcgggca agttggctga gggcgtgcgg 60
cagaggctgc thccctcggc gacgcgaccc ctcagcaact caagctatga actgaagctc 120
cctagggacg gagaccggag cggagcggcg gaggcagcag cagcagcagc agcagcagca 180
gcagcagcag ccgccgccgc cgcgcctta gcgggaactg agcagaccg gcgcggagcc 240
acgactcctg cacgthtacc thctgtcgc cgttctctgc ggcggttgcc taaaagacgt 300
tacagccgc agaccgaca caccaaagcc gctthtctcc gcgccccgc ccaggggaggc 360
tgccggccag aagggacccc acctgagagc agctcgggct gctgagttcg thttgtgtct 420
gagctctgcg ctctgcacgg aaccgacccc gtacccatgg ctctgataat ggaaccggtg 480
agcaaatggt ctccgagtca agtagtgagc tggatgaaag gtcttgatga ctgtthgcag 540
cagtatatta agaactthga gagggagaag atcagtgagg accagctgct gcgcattaca 600
catcaggagc tagaagatct gggggctcag cgcattggcc atcaggaact gatcttgga 660
gcagttgacc thctgtgtgc attgaattat ggcttggaag cagaaaatct aaaaacctt 720

```

tctcacaagt	tgaatgcac	tgccaaaaat	ctgcagaatt	ttataacagg	aaggagaagg	780
agtggccatt	atgatgggag	gaccagccga	aaattgccaa	acgactttct	gacctcagtt	840
gtggatctga	ttggagcagc	caagagtcctg	cttgccctgg	tggacagggtc	accattttgct	900
gctgtgacag	actattcagt	tacaagaaat	aatgtcatat	aactctgcct	ggagtttaaca	960
acaattgtgc	aacaggattg	tactgtatat	gaaacagaga	ataaaattct	tcacgtgtgt	1020
aaaactcttt	ctggagtctg	tgaccacatc	atatccctgt	cgtcagatcc	tctgggtttca	1080
cagtctgctc	acctggaagt	gattcagctg	gcaaacatta	aaccaagcga	agggctgggt	1140
atgtatatta	aatctacata	tgatggcctc	catgtaatta	ctggaaccac	agaaaattca	1200
cctgcagatc	ggtgcaagaa	aatccatgct	ggcgatgaag	tgattcaagt	taatcatcag	1260
actgtgggtg	ggtggcagtt	gaaaaatttg	gtgaatgcac	tacgagagga	cccagtggt	1320
gttatcttaa	ctttgaaaaa	gcgacctcag	agcatgctta	cctcagcacc	agctttactg	1380
aaaaatatga	gatggaagcc	ccttgctctg	cagcctctta	tacctagaag	ttcccacaagc	1440
agcgttgcca	cgccttcag	caccatcagt	acacccacca	aaagagacag	ttctgccctc	1500
caggatctct	acattcccc	tcctcctgca	gaaccatata	ttcccaggga	tgaaaaagga	1560
aaccttcctt	gtgaagacct	cagaggacat	atgggtgggca	agccagtgc	taagggatct	1620
gaatcaccaa	attcattttct	ggatcaggaa	tatcgaaaga	gatttaatat	tgtcgaagaa	1680
gatactgtct	tatatgtcta	tgaatatgaa	aaaggaagat	caagtagtca	aggaagacga	1740
gaaagcacc	caacttatgg	caagctacga	cctatatcta	tgccagtgg	atataattgg	1800
gtgggggact	atgaagatcc	aaataagatg	aagagagata	gtagaagaga	aaactctcta	1860
cttcggtata	tgagcaatga	aaagattgct	caagaagaat	acatgtttca	gagaaacagc	1920
aaaaaggaca	caggggaagaa	gtcaaaaaag	aaggggtgata	agagtaatag	cccaactcac	1980
tattcattgc	tacctagttt	acaaatggat	gcactgagac	aagacatcat	gggcactcct	2040
gtgccagaga	ccacactata	ccatacattt	cagcagtcct	cactgcagca	caaatcaag	2100
aagaaaaaca	aaggtcctat	agcaggcaag	agcaaaagac	gaatttcttg	caaagatctt	2160
ggcctgtgtg	actgtgaggg	ctggccttgg	aaaaagaaag	atgcgaagag	ttacttttca	2220
cagaaatgga	aaaaatattg	gtttgtccta	aaggatgcat	ccctttattg	gtatattaat	2280
gaggaggatg	aaaaagcaga	aggattcatt	gcctgtcctg	aattttaa	tgatagagcc	2340
agtgaatgcc	gcaaaaaata	tgcatctcaa	agctgtcatc	ctaaaatcaa	aagcttttat	2400
tttgcctgtg	aacatcttga	tgatatgaac	aggtggctta	acagaattaa	tatgctgact	2460
gcaggatatg	cagaaagaga	gaggattaag	caggaacaag	attactggag	tgagagtgc	2520
aaggaagaag	cagatactcc	atcaacacca	aaacaagata	gccctccacc	cccatatgat	2580
acataccac	gacctccctc	gatgagttgc	gccagtccct	atgtggaagc	aaaacatagc	2640
cgactttcct	ccacggagac	ttctcagctc	cagtcctctc	atgaggagtt	tcgccaggaa	2700
gtaactggga	gcagtgcagt	gtctccatt	cgcaagacag	ccagtcagcg	ccgctcctgg	2760
caggatttaa	ttgagacgcc	actgacaagt	tcaggcttac	actatcttca	gactctgccc	2820
ctggaggatt	ctgtcttctc	tgactccgcg	gccatctccc	cagagcacag	gcggcagctc	2880
accctgccaa	ctcagaaatg	ccacctgcag	gatcactatg	ggccataccc	cttagctgag	2940
agtgagagga	tgcaagtgc	aaatggaaat	gggggcaagc	ctcgaagttt	tactctgcct	3000
cgagatagcg	ggttcaacca	ttgctgtctg	aatgctccag	ttagtgcctg	tgaccacag	3060
gatgacgtgc	aacccccaga	ggtggaggaa	gaggaggagg	aggaggagga	ggaaggggag	3120
gcagcagggg	aaaacatagg	agaaaaagc	taatacactg	cgagagttgg	tagaacctct	3180
ccatgccaaa	tcggatccac	ttctgttggc	actcaaccca	ttggactcac	agattgataa	3240
gctaattgtt	agagaattta	gatcggagag	agtcggtacg	gcgcagactc	aacatcaacc	3300
tcttgcaagc	aactaaaatg	gcctcgtcct	tgctgtttat	aaagggatat	agacttgtaa	3360
aaagcttaga	tcatacaagt	ttttggattg	ggggcctccc	acattgggac	aagaggggca	3420
ggccactcct	aagaagaatg	cgactttctc	acattgggac	tagcataaga	tcaaagccaa	3480
tcagatgga	gcacagtaac	agaaaaactgc	ggtttctgtg	ggagaacaga	aggggaaagg	3540
gtcttaactg	ggaaagggct	ctgtgtggta	acacctcagt	tgtgttctcc	tgacaccagg	3600
aaaagagagg	gatcagcttc	aataactaga	aaattctggc	tgtttaatgg	actctttggg	3660
ggcctcttta	aggcaaagca	gagaaagcaa	attatgtatt	aagtgtattt	tgcattttta	3720
aaacttgacg	tgctgtattg	tactaaatta	agtgtaatct	attaaggcaa	ggtatacaca	3780
atgtgctttg	aaacttacta	tgtttattct	attataaagt	gtattcaggt	gcaacacaga	3840
gactgctttc	ggtgacatta	atgaagaaaa	tttctcatgc	caggctttat	tatagaatct	3900
tcagctaaaa	tcctaacttt	ctccttattt	cttggcactt	gtatacaagt	ggtgttgcct	3960

cttagggcag gcatgagcta ttcttttctg taaaatattt tgaatctata ggctgtgggt 4020
ttcatttttg aaaagtattt tgtctggatg tctttcaaac tagcttcaga tattatttaa 4080
tactatgtaa ctgggtcccc tatgggtcaa tcaatattgc ttatttttct tctgtagtgg 4140
atgtgaaatt tccttttagt ggataagata cactgtaata attttaatgc taattaatga 4200
tatttcatac tgtgcaatga acagataatt taacactgta ttttgaaatg ttttttctt 4260
cctgtcaccg cagtgtgtgg tattgcataa tgtgaatacc tgtaaaaata taaattactt 4320
aaaaataaaa atatgaccaa ttggtatcag 4350

<210> 7

<211> 50

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA fragment
of Maguin-1

<400> 7

ggagagaggc agcactgtaa actgaagtca aataaattca gctcttaatg 50

<210> 8

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
One-base-anchor oligonucleotides

<400> 8

httttttttt tta 13

<210> 9

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
One-base-anchor oligonucleotides

<400> 9

httttttttt ttg 13

<210> 10

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
One-base-anchor oligonucleotides

<400> 10
htttttttttt ttc

13

<210> 11
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for
Maguin-1

<400> 11
cagcaagcag ttgacggga

19

<210> 12
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for
Maguin-1

<400> 12
tgaatttgac aagactcgtg gc

22

<210> 13
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for
Maguin-2

<400> 13
gggcctccca aagggatat

19

<210> 14
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for
Maguin-2

<400> 14
cccaatgtag aaagctcgca tt

22

<210> 15
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer for
cyclophilin B

<400> 15
actgaagcac tacgggcctg

20

<210> 16
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer for
cyclophilin B

<400> 16
agccgttggt gtctttgcc

19

<210> 17
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer for
ribosomal protein S9

<400> 17
gggtcaaattt accctggcca

20

<210> 18
<211> 22
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer for
ribosomal protein S9

<400> 18

tctcatcaag cgtcagcagt tc

22

<210> 19

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer for
beta-actin

<400> 19

tggaacgggtg aaggtgaca

19

<210> 20

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer for
beta-actin

<400> 20

ggcaagggac ttcctgtaa

19

<210> 21

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer for
GAPDH

<400> 21

cgtcatgggt gtgaaccatg

20

<210> 22

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer for
GAPDH

<400> 22

gctaagcagt tggtagtgca g

21

<210> 23
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for
transferrin receptor

<400> 23
gtcgctggtc agttcgtgat t 21

<210> 24
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer for
transferrin receptor

<400> 24
agcagttggc tgttgtacct ctc 23

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 03/02857

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/47 C12N15/12 C12N5/10 C07K16/18 G01N33/53
C12Q1/68 A61K38/16 A61K48/00 A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K G01N C12Q A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

GENSEQ, EMBL, EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! 16 January 2002 (2002-01-16) LANIGAN, T.M. AND GUAN, K.L.: "Homo sapiens connector enhancer of KSR2A mRNA, complete cds." Database accession no. AF418269 XP002210679</p> <p style="text-align: center;">--- -/--</p>	<p>11, 13, 14, 16, 20, 26, 28, 29</p>

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

18 June 2003

Date of mailing of the international search report

30/06/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

ALCONADA RODRIG., A

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 03/02857

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE GENSEQ 'Online! 8 November 2000 (2000-11-08) TAKAI, Y. ET AL.: "Rat MAGUIN 1 protein" Database accession no. AAY92942 XP002210736 -& DATABASE WPI Section Ch, Week 200033 Derwent Publications Ltd., London, GB; Class B04, AN 2000-387785 XP002210680 & WO 00 29572 A (KAGAKU GIJUTSU SHINKO JIGYODAN), 25 May 2000 (2000-05-25) abstract</p> <p>---</p>	<p>11,13, 14,16, 20,26, 28,29</p>
X	<p>DATABASE GENSEQ 'Online! 8 November 2000 (2000-11-08) TAKAI, Y. ET AL.: "Rat MAGUIN 2 protein" Database accession no. AAY92943 XP002210737 -& DATABASE WPI Section Ch, Week 200033 Derwent Publications Ltd., London, GB; Class B04, AN 2000-387785 XP002210680 & WO 00 29572 A (KAGAKU GIJUTSU SHINKO JIGYODAN), 25 May 2000 (2000-05-25) abstract</p> <p>-----</p>	<p>11,13, 14,16, 20,26, 28,29</p>

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 12-15 (in part) and 19, 21-25 (complete)

Present claim 12 relates to a method of treating or preventing a neurodegenerative disease, said method comprising administering to said subject an amount of an agent, wherein said agent is defined by reference to a desirable characteristic or property, namely, that said agent directly or indirectly modulates the activity or level of the (i) gene coding for human MAGUIN-1 and/or human MAGUIN-2; and/or (ii) a transcription product of a gene coding for human MAGUIN-1 or human MAGUIN-2 and/or (iii) a translation product of a gene coding for human MAGUIN-1 and/or human MAGUIN-2 and/or a fragment, or derivative, or variant of (i) to (iii). The claims cover the methods that involve the administration of an agent having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT for only a very limited number of such agents. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the agent by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the methods which make use of the agent, wherein the agent comprises the human MAGUIN-1 or MAGUIN-2 polypeptide sequences or the corresponding polynucleotide sequence in sense or antisense orientation (see page 14 from the description).

The same objection applies to claim 13, which refers to the above mentioned agent or modulator as such, to claim 14 which refers to pharmaceutical compositions comprising said modulator and to claim 15, which refers to the second medical use of said modulator.

Present claim 19 relates to a method for the identification of a compound for inhibition of binding between a ligand and human MAGUIN-1 or MAGUIN-2, wherein said method implies contacting MAGUIN-1 or MAGUIN-2 with a detectable ligand in the presence of the compound to be tested. The claim covers all possible MAGUIN ligands, whereas the application does not provide support within the meaning of Article 6 PCT or disclosure within the meaning of Article 5 PCT for any of such MAGUIN ligands. In the present case, the claim so lacks support, and the application so lacks disclosure, that a meaningful search of the claim is impossible. Independent of the above reasoning, the claim also lacks clarity (Article 6 PCT). An attempt is made to define the product by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been not carried out for those claims.

Claims 21-25 relate to methods of producing medicaments and medicaments

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

obtained by said method, wherein said medicaments contain a compound which is defined by reference to a desirable characteristic or property, namely, that the compound can be identified by the methods of claim 18-20. The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT for none of such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search of the claim impossible. Consequently, the search has not been carried out for those claims.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/EP 03/02857

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 12 and 17 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 12-15 (in part) and 19, 21-25 (complete)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.